



CHIRONOMUS Newsletter on Chironomidae Research

No. 22

ISSN 0172-1941

October 2009



Yin and yang? *Chironomus dilutus* gynandromorph. Photo: Quentin Lang, Department of Genetics, The University of Melbourne

CONTENTS

Editorial: XVII International Symposium on Chironomidae, July 5-10, 2009, Tianjin, China.....	3
Trends and development in chironomid palaeoecology: Summary from the 9th workshop on subfossil Chironomidae.....	5
Memories from the XVIIth chironomid symposium in Tianjin, China.....	8
Chironomid research in northern Patagonia.....	9
Development of a new design of an insectary model for rearing and environmental assessment studies on Chironomid midges.....	11
Current Research	14
Martin, J. & Saxena, S.: Synonymy of <i>Chironomus plumatisetiger</i> Tokunaga, 1964, with <i>Chironomus circumdatus</i> Kieffer, 1916.....	14
Paggi, A.C., Siri, A. & de Oliveira, C.S.N.: A brief analysis of morphology and terminology for the genus <i>Ablabesmyia</i> ..	15
Agius, J.T., Gibson, J.A.E, Rico, E. & Quesada, A.: Paleolimnological evidence confirms that <i>Parochlus steinenii</i> (Gerke) is not a recent introduction to the Antarctic Peninsula Region.....	18
Short communications	22
Current bibliography	27

CHIRONOMUS Newsletter on Chironomidae Research

Editors

Torbjørn EKREM, Museum of Natural History and Archaeology, Norwegian University of Science and Technology, NO-7491 Trondheim, Norway

Peter H. LANGTON, 5 Kylebeg Avenue, Mountsandel, Coleraine, Co. Londonderry, Northern Ireland, BT52 1 JN, Northern Ireland

Current Bibliography

Odwin HOFFRICHTER, Institut für Biologie I, Albert-Ludwigs-Universität Freiburg, Hauptstrasse 1, D-79104, Germany

Treasurer

Trond ANDERSEN, Natural History Collections, Bergen Museum, University of Bergen, Musépllass 3, NO-5007 Bergen, Norway

CHIRONOMUS Newsletter on Chironomidae Research (ISSN 0172-1941) is published yearly in October. Submission deadline for contributions is July 1.

Contributions to *CHIRONOMUS Newsletter on Chironomidae Research* should be submitted per e-mail to: Torbjørn Ekrem: Torbjorn.Ekrem@vm.ntnu.no or Peter H. Langton: PHLangton@kylebegave.fsnet.co.uk. Please use the following formatting: Text in 12 point Times New Roman, first page must include title, name, address and email address of all authors. Headings should be bold faced. Cite relevant references in parentheses without comma between name and year [ex. (Langton 1991)]. List all references alphabetically in the format of the *Current Bibliography* at the end of the manuscript. Tables can be included directly in the text. Text should preferably be submitted as MS Word or rtf files. All figures should be supplied separately as tiff or jpg files.

Would you like to see your picture on the front page? Please send us your favourite midge photograph or drawing (Torbjorn.Ekrem@vm.ntnu.no).

Editorial

XVII International Symposium on Chironomidae, July 5-10, 2009, Tianjin, China

It must be the dream of many people in the world to visit China and for us this dream came true: China is awesome and the XVIIth International Symposium on Chironomidae was a real joy to attend. Our hosts, Xinhua Wang, Wei Liu and their student helpers turned themselves upside down to make us at our ease and have the program run smoothly. Nothing seemed too much for them: they were the perfect hosts.

The Symposium began on the Sunday evening with a social gathering that brought together long standing researchers and students in the Chironomidae in a relaxed and convivial environment. A very enjoyable event with 60+ delegates from every biogeographical region except the poles - a lot to catch up on and new contacts to make. It was unfortunate that the distances involved and the recession meant that some previous symposium participants were unable to attend, but this venue opened up the conference to far eastern colleagues, so fulfilling one of the Symposia's roles of gathering together, in dialogue, chironomid researchers from across the world.

The scientific programme started at 9.00 the following morning in a spacious new lecture theatre in the new Building of the College of Life Sciences with a short inauguration ceremony, followed by the Thienemann Honorary Lecture, an excellent and polished delivery by Jon Martin entitled "From bands to base pairs: Problems in the identification of *Chironomus* species, using the example of *Chironomus oppositus* Walker." Then the series of oral communications commenced, 29 papers authored and co-authored by about 60 researchers. Papers covered morphology, molecular phylogeny, karyology, systematics, ecology and biogeography; equally diverse were the ages of those giving the talks: from 16 to 70+! An outstanding array of posters embellished the walls of the large airy foyer to the lecture theatre, where delegates browsed and discussed the submissions at break times while consuming many lychees! The program and abstracts can be viewed on the symposium website: <http://entomology.nankai.edu.cn/17chiro>. For one (Peter) who has been studying these insects for half a century, there was much to marvel at and a question to ponder: will the use of DNA sequences to clarify relationships and delineate species reduce the opportunities for biologists without such facilities/techniques to contribute to progress in the Chironomidae?

The Symposium banquet was held in an astonishing restaurant: a massive greenhouse containing a tropical jungle of mature trees and palms under which the tables were distributed. In the centre of each circular table was a turntable, to which were added time and time again new dishes to try: a really exotic gustatory experience in an equally exotic environment. Through this 'jungle' flows a stream populated by large Koi Carp and a lagoon that contains seals which add a touch of the incredulous to the already fantastic venue.

On the Wednesday we were taken on a river boat cruise of the Hai He river that runs through Tianjin. Although the skim net was in the water for the duration, no exuviae were obtained, presumably due to the mid summer pause in emergence. This was a great way to see outstanding examples of the architecture of modern Tianjin and of the many

different highly ornate bridges under which we passed. We visited an exceedingly modern museum with exceptional ambience containing artefacts from distant history to the modern day; the early artefacts were especially fascinating as China has such a long cultural history.

The Symposium concluded with a forum session on Thursday afternoon: the next symposium in two years' time will be in Trondheim, courtesy of Torbjørn Ekrem and Elisabeth Stur. Their presentation gave the delegates a taste of the area – so different from that of the present venue! – and their proposed arrangements.

The post symposium tour took place on the Friday. A coach took us to experience the Great Wall, with views of the wall receding into the distance perched on the ridges of successive hills. It is indeed a marvel of human endeavour and we felt privileged to be walking a portion of it. After that we were transported into Beijing and were guided through the Forbidden City in reverse – a concession to parking the coach! The succession of halls and courtyards were awe inspiring and a real insight into China history.

Amongst our souvenirs we always have some entomological trophies: this time we have two large cicadas (the song of cicadas filled the air wherever we went all day – a similar sound took over at night generated by presumed bushcrickets), some cicada exuviae, two large prettily coloured homopteran bugs and one chironomid pupal exuviae collected in a lotus filled canal at the university: *Glyptotendipes paripes*!!

Xinhua Wang, Wei Liu and their helpers are to be congratulated on a most impressive and very well organised symposium and providing great memories through food and visits. On behalf of all participants we propose a big THANKYOU!

Peter Langton and Sue McBean, 5 Kylebeg Avenue, Mountsandel, Coleraine, Co Londonderry, BT52 1 JN, Northern Ireland. E-mail: PHLangton@kylebegave.fsnet.co.uk.

TRENDS AND DEVELOPMENT IN CHIRONOMID PALAEOECOLOGY: SUMMARY FROM THE 9TH WORKSHOP ON SUBFOSSIL CHIRONOMIDAE

Klaus Peter Brodersen¹, Gaute Velle² & Nina S. Reuss¹

¹*Freshwater Biological Laboratory, Biological Institute, University of Copenhagen, Denmark.*
kpbrodersen@bio.ku.dk, nreuss@bio.ku.dk

²*Bergen Museum, University of Bergen, Norway.* nzlgv@uib.no

The 9th workshop on Subfossil Chironomidae was held at the Biological Institute, University of Copenhagen on May 4–6, 2009. There were 27 palaeo-chironomid researchers present, representing 14 different nationalities (Fig. 1, group photo). The purpose with these workshops is to exchange ideas and to have critical and fruitful discussions about topics ranging from larval identification to ecological interpretation – whether people are working with subfossil or contemporary data. Problems and progresses in chironomid palaeoecology are discussed among people with their fingers deep in the mud, and the informal presentations give a fine overview of current state-of-the-art research and development.

The workshop was organised into six sessions with oral presentations, one session of general discussion and a one-day field excursion. The field excursion on the first day took us around to the North Zealand birch forest in full leaf. We had a short lunch break at the Freshwater Biological Laboratory belonging to the University of Copenhagen in the town of Hillerød. At the shore of Lake Esrum, the laboratory's Grand-Old-Man,

Pétur Jónasson (born 1920) (Fig. 2), gave a fine summary of his illustrious work on life-cycles, respiration, secondary production and benthic-pelagic coupling of *Chironomus anthracinus* in Lake Esrum (Jónasson 1972).

With three sessions under the climate topic, it is evident that chironomid-inferred temperature reconstruction is still a central issue. Subfossil chironomids are still used as the primary quantitative climate indicator in multi-proxy studies of e.g., Early Weichselian, as presented by Stefan Engels, and Late Glacial sediments, as presented by Mateusz Plociennik. New quantitative temperature models are being developed and improved. Results of two merged European data sets were presented by Steve Brooks, a Finnish data set was shown by Tomi Luoto, and two independent data sets from Northern Russia are being developed and were presented by Larisa Nazarova and Angela Self. Along with these studies, preliminary stratigraphies and taxonomic problems were addressed by Verana de Hoog. A novel idea from Angela Self was to incorporate “continentiality” into



Figure 1. Participants at the workshop. Photo: Katrine Kongshavn.

the quantitative models. Down-core continentality reconstructions from the Putoran Plateau suggest a lengthening of the growing season despite relatively constant temperature over the 1950s–2000.



Figure 2. Professor Pétur M. Jónasson speaking about *Chironomus* population dynamics at the shore of Lake Esrum. Photo: Katrine Kongshavn.

Analyses of recent climate change over the last few hundred years were presented from the far north in Spitsbergen (Katrine Kongshavn) to the south in Patagonia (Peter Langdon). Comparisons to instrumental and independent records (e.g., known past fluctuations in glacier extent), are used in order to assess the potential for reconstructing accurately climate change over longer (millennial scale) time periods (Peter Langdon). High-resolution analyses from annually varved sediments are likewise used along with instrumental records in evaluation of model performance in Switzerland (Isabelle Larocque).

Studies of past environmental change in the Bolivian Andes were presented by Joseph J. Williams. In this project they are tracing the natural history of an endangered woodland ecosystem in order to evaluate if the patchy distribution of *Polylepis* observed today is a relict of past more widespread dominance and if the current distribution is human and/or climate driven. The chironomids are used as independent climate proxies.

Human induced environmental impacts, aside from climate change, were in focus of many studies. Enrichment (eutrophication) effects of human settlement were discussed from Northern Ireland by Naomi Holmes. Models across Europe have suggested that events such as the adoption of agriculture in Europe are climatically driven and hints of this relationship are present in existing North Mayo data. The environmentally sensitive nature of the area means that North Mayo is an ideal case study for the relationships between climate change and changes in human settlement.

From Africa, Bob Rumes demonstrated a clear evidence for anthropogenic presence and influence on the landscape in Western Uganda dating back 2000 years. Although there were pronounced natural climate variations in the area, the effect of these is not always clear and, at times, difficult to distinguish from anthropogenic changes. Complex ecological responses to eutrophication and recent climate in Denmark were analysed against environmental monitoring data using fine interval sampling (Wing Wai Sung). Recent human impact on lake food webs by e.g., introduction of fish is also a new field in chironomid palaeolimnology. Such studies are carried out in subalpine lakes in France (Virgile Baudrot), in south Swedish lakes (Kimmo Tolonen) and in crater lakes at the Azores (Klaus P. Brodersen). The role of natural changes, such as succession in primary producers, food resources, oxygen conditions, intra- and inter-specific competition and changes in benthic communities is studied by Nina Reuss. She is studying stable isotope signatures on the most common chironomid taxa from West Greenland as well as on their potential food resources. Successful results will allow down-core inference of changed food-web structures and lake dynamics.

Examination of fossil midge assemblages from surface sediments of small shallow lakes in Finland took us back to basics in interpretation of chironomid palaeolimnology. Tomi Luoto presented multi- and intra-lake studies to emphasize that the controlling environmental factors are scale dependent from temperature over lake depth to hypolimnetic oxygen conditions. Intra-lake dataset from eastern Finland showed that the most important environmental factors controlling midge distribution within a lake basin were river contribution, water depth and submerged vegetation patterns. It also appeared that the fossil midge assemblages represented fauna that lived in close proximity to the sampling sites thus enabling investigation of intra-lake gradients in midge assemblages.

Novel studies on the potential of using subfossil chironomids as indicators of palaeoflow regimes were discussed by Lynda Howard. Chironomidae, identified to species level where possible, could be classified into groups of “Lotic invertebrate Index for Flow Evaluation”. This information was used to reconstruct the palaeoflow regime of a river and associated changes to in-stream habitats.

Improvement of subfossil taxonomy and identification has always been a central subject in these workshops. Boris Ilyashuk presented

current knowledge and palaeoecological perspectives regarding the genus *Pseudodiamesa*. The aim was to clarify the morphological characters of head capsules of the *Pseudodiamesa* larvae which can be clearly visible within subfossil remains and reliable used for an identification of the subgeneric morphotypes. A short summary of the zoogeography and ecology of *Pseudodiamesa* species was also given which is very helpful in palaeoecological interpretation of subfossil records.

A general discussion on palaeoenvironmental inferences at the extremes of the environmental gradient was introduced and chaired by Gaute Velle. The modern calibration data set lake samples include chironomid head capsules and concurring environmental variables. However, it is not straightforward to use the calibration data set to infer environmental variables that are near the lower or upper end of the gradient captured by the modern data. Gaute Velle pointed out typical challenges, such as:

- (1) The edge effect. Inferring variables at the end of the gradient must be done with caution. The so-called 'edge effect' caused by a truncation of the taxa responses, results in a tendency to pull the predicted values towards the mean of the environmental range in the calibration data set.
- (2) Few modern taxa at the extremes. Since there often are more lakes in the mid-range of the calibration data sets, fewer taxa are captured from the gradient extremes. Hence, the response functions for the taxa at the extremes are less confident as they are based on fewer observations.
- (3) Low fossil diversity. Sites or time periods with extreme environments may have additional challenges as these often have a low diversity. When there is a low diversity in the fossil assemblage, the taxa that are present influence the palaeoenvironmental inference to a large extent and the inference may be more prone to bias caused by inadequate response functions or to stochastic variables.
- (4) Low abundance. Sites or time periods at extreme environments often have a low abundance of chironomids. Any quantitative palaeoenvironmental inference should be based on >50 head capsules.
- (5) Pioneer assemblages. In extreme habitats, such as newly formed cold proglacial lakes, the

pioneer fauna may be in disproportion with concurrent environments because they are specialist pioneers or because they are opportunists.

In summary, the workshop was very fruitful and inspiring. Chironomid palaeoecology is still growing, but also developing away from blind routine temperature reconstructions of past climate conditions. In the summary of the previous workshop in Reykjavik, Langdon *et al.* (2007) stated: "...in other words palaeoecological simplicity may need to be considered more carefully in the face of ecological complexity". Many discussions at the workshop clearly demonstrated that other human and natural impacts, aside from climate and temperature, are affecting the fossil assemblages, and that many of these parameters can be successfully interpreted. It is also positive that many from the new generation of chironomid palaeoecologists are going back to basics to critically assess the strengths and weaknesses of their models and methods. We strongly encourage such efforts and also efforts made to integrate neo- and paleolimnological approaches to refine interpretations of environmental change (e.g., Saros 2009). Such efforts are important in order to understand the biological mechanisms driving the species distribution and abundance.

We thank all for active participation and constructive discussions. The next workshop on subfossil chironomidae is arranged by Gaute Velle in Bergen, Norway, in May 2011.

References

- Jónasson, P.M. 1972. Ecology and production of profundal benthos in relation to phytoplankton in Lake Esrom. - *Oikos Supplementum* 14: 1–148.
- Langdon, P., Holmes, N., Stefánsson, S.M., Hannesdóttir, E. and Axford, Y. 2007. The 8th European Subfossil Chironomid Workshop, Reykjavik 7–8th May 2007. - *Chironomus newsletter on Chironomidae research* 20: 10–13.
- Saros, J. 2009. Integrating neo- and paleolimnological approaches to refine interpretations of environmental change. - *Journal of Paleolimnology* 41: 243–252.

MEMORIES OF THE XVIITH CHIRONOMID SYMPOSIUM IN TIANJIN, CHINA

This was the first time I have attended an International Symposium on Chironomidae, held in Tianjin, China, during July 5-10, 2009. It was an excellent way to contact people working in the area and was a great experience to discuss their work on chironomids. I was very well integrated in the group from the first day and I strongly recommend this Symposium to all people working with Chironomidae. See you in Norway!

Pedro Raposeiro, PhD Student, Universidade dos Açores discute, Departamento de Biologia, Rua Mãe de Deus, 13A 9501-801 Ponta Delgada, Portugal.

My name is Mónika Tóth from the University of Debrecen, Hungary. This was the second time that I have attended the International Symposium on Chironomidae. I think this conference was a good opportunity for students and Ph.D. students, who deal with non-biting midges, to meet with academic scientists and ask for help with their work. This time I had a poster and I got valuable recommendations and comments to continue my studies. I enjoyed the conference and interesting conversations. This was my first time in China, which is a very amazing country. Thank you Xinhua Wang, Liu Wei and their students for the organisation of the symposium.

Mónika Tóth, University of Debrecen, Hungary.

The Tianjin Symposium was a fantastic introduction to Chironomidae Symposiums and certainly what I hope will be the first of many. The 'seasoned' delegates provided a very welcoming environment for 'first timers', and the enthusiasm of all attendees was quite stimulating. I found the quality of the presented research to be phenomenal, and the Symposium was the perfect venue to have exciting conversations with

scientists from around the world who I had known before only through their publications. Additionally, having the conference in Tianjin provided many of us with the opportunity to broaden our cultural horizons as well as scientific: experiences such as a trip to the Great Wall, a river boat trip on the Hai He River, trying out traditional Chinese cuisine, and, of course, having ample opportunity to find the perfect souvenir provided cultural treats that could please everyone. All-in-all, the symposium was very well organised, a great opportunity to make new contacts, and an opportunity to learn from other cultures. I'm already looking forward to the next conference in Trondheim!

Alyssa M. Anderson, Department of Zoology, University of Minnesota, St. Paul, Minnesota 55108.

The 17th International Symposium on Chironomidae was held in Nankai University this year. It was a distinct opportunity for us to meet a number of renowned experts working on different fields of Chironomidae, meanwhile updated information was conveyed. The lectures they delivered regaled our senses and helped quite a lot in our study. The posters that the participants brought showed their research results and created a kind of bright scenery in the hall of the meeting place.

Being service staff, we felt tired but excited and delighted after each busy day. The smiling faces of the foreign friends will not be forgotten till the cows come home. Both of us wish the close friendship to grow deeper and deeper between the researchers all over the world and Nankai University through this symposium. What's more, we are looking forward to the next symposium where we can get together again.

Hui Sun & Xue Yu, College of Life Sciences, Nankai University.

CHIRONOMID RESEARCH IN NORTHERN PATAGONIA

Donato, Mariano¹, Julieta Massaferrero² & Stephen J. Brooks³

¹*Laboratorio de Sistemática y Biología Evolutiva (LASBE), Museo de La Plata, Paseo del Bosque s/n (1900), La Plata, Argentina, mdonato@fcnym.unlp.edu.ar*

²*Laboratorio de Biodiversidad, INIBIOMA - CONICET, San Martín 24, (8400) Bariloche, Argentina, julimassaferrero@crub.uncoma.edu.ar*

³*Department of Entomology, Natural History Museum, London SW7 5BD, United Kingdom, sjb@nhm.ac.uk*

Patagonia is the southernmost territory of South America and was traditionally considered as belonging to the Neotropical region (Sclater 1858, Wallace 1876). Based on these zoogeographic schemes, the Argentinean biogeographer Raul Ringuelet (1961) divided Patagonia into the Araucanian Subregion, characterized by dominance of Austral or Notogeic fauna, and the Andean-Patagonian Subregion, having a dominant mixture of Notogeic, Brasilic, and Nearctic elements. Later, Cabrera & Willink (1973) proposed a biogeographic scheme for South America and the Caribbean in which the Andean-Patagonian region was part of the Neotropical region and Araucania. They named the region the Sub-Antarctic province, which was part of the Antarctic region and was based on the floristic and faunistic affinities of southwestern Patagonia with Antarctica, Australia and New Zealand. Further biogeographic schemes considered Patagonia as a part of a Holantarctic region (Takhtajan 1986) or Sub-Antarctic subregion and Patagonian subregion of the Andean Region (Morrone 2001). The extraordinary amount of different biogeographic schemes reflects the complexity of the affinities of the biota of Patagonia.

From an ecological perspective, Patagonia is highly influenced by the Andes cordillera. The climate in western Patagonia is cool-temperate but rainfall decreases drastically eastwards towards the steppes. Therefore, climate and soil conditions make Patagonia extremely water-limited and this is reflected in the aquatic ecosystems. The western part of Patagonia is covered by two eco-regions. The Valdivian temperate forest, covering a narrow continental strip between the western slope of the Andes and the Pacific Ocean, and running from 35°S to 48°S, is characterized mainly by deciduous and evergreen forests and Valdivian rainforest. The eastern part of Patagonia is distinguished (from north to south) by the Argentinean Monte, dominated by scrub vegetation, the Patagonian steppe, with shrub vegetation, and the Patagonian

grassland, with grass-steppe interspersed with shrubs (Olsen et al. 2001).

The habitats and communities in Patagonian freshwater ecosystems have been modified by different extents and causes. The construction of dams and the resulting reservoirs to regulate rivers and control floods have modified seasonal flow patterns, thus affecting environmental conditions for most species. As a result, lotic watercourses have become lentic environments. Other important alterations of the physical environment in Patagonia are the consequence of mining activities, urban development and degradation derived from tourism. Ironically, wetlands are essential for sustaining biodiversity and wildlife. Patagonian wetlands present complex aquatic communities and provide habitat for threatened species and a high number of endemic species.

The freshwater insect biodiversity of Patagonia is poorly known and knowledge is constrained by a lack of adequate identification guides and reference collections. In addition, there is poor public understanding of the importance of wetlands for biodiversity. In order to address these issues, a three year Darwin Initiative project, entitled "Capacity Building for Aquatic Insects in Patagonia" and supported by the Department for Environment, Food and Rural Affairs (Defra, UK), began in 2006. The study area is the Nahuel Huapi National Park (NHNP) in northern Patagonia, Argentina. NHNP covers 7050 km² of Patagonian steppe, temperate deciduous forest and Valdivian rainforest. The latter ecosystem is recognised as one of the most threatened eco-regions in the world by the Global 2000 initiative launched by WWF and the World Bank. The main objectives of the Darwin Initiative project are to increase knowledge of freshwater insect biodiversity; to promote conservation of the various freshwater ecosystems in this protected area; and to communicate, share and educate citizens in different aspects of biodiversity. The project brings together a team of British and Argentinean scientists from the Natural History Museum

London, The Natural History Museum La Plata, The Institute of Limnology La Plata, The University of Patagonia in Esquel, The Central Regional University at Bariloche and the National Parks Administration Bariloche.

A key task for the team is to study the major freshwater insect groups in NHNP including Chironomidae, Ceratopogonidae and Simuliidae (Diptera); Plecoptera; Ephemeroptera; Odonata; Trichoptera; and aquatic Coleoptera. With invaluable logistics support from the park rangers and other NHNP staff, more than 200 sites have been sampled covering the various freshwater environments represented in NHNP. The sampling methods included Malaise trap, light trap, sweep net, drift net and kick sample. Taxonomic, ecological and distributional data resulting from this collecting effort is being entered into a database of freshwater insects from NHNP. The species distribution data will be linked to a vegetation classification and the physical and chemical attributes of the water bodies, using GIS and digital imaging, and will be used to model freshwater insect data spatially and create a biodiversity data repository, the first of its kind in Patagonia. Another important project goal was addressed by building infrastructure in NHNP to provide a wetland interpretation centre and laboratory where tourists, sport anglers, students and researchers can study freshwater insects and understand their role in freshwater ecosystems.

Patagonian chironomid fauna

The first records and knowledge of Chironomidae from Patagonia are due to the invaluable work of the English entomologist Frederick Wallace Edwards (1888-1940). To obtaining collections of insects from the southern Andes, a joint expedition was arranged in 1926 by the British Museum of Natural History and the Bacteriological Institute of the National Department of Hygiene of Argentina. Edwards went to collect for the British Museum and Mr. Raymond C. Shannon for the Argentine Government. At the end of the field trip, they collected collections of over 20000 insects each one (Edwards 1927). After the study of the material collected, Edwards published in 1931 published the chapter referred to Chironomidae as a part of the book "Diptera of Patagonia and South Chile" and he described 71 new species and gives the foundation of the chironomid Patagonian knowledge.

Preliminary data on Chironomidae collected during the project, supplemented by examining

collections made by earlier entomologists and literature searches, has shown that the chironomid fauna of NHNP includes 104 species in 48 genera and 6 subfamilies. Analysis of subfossil larvae from lake sediments collected in NHNP include 52 morphotypes in 36 genera and 4 subfamilies (Donato et al. 2008). The number of chironomid species found in NHNP constitutes a relatively high proportion of the chironomid diversity of Patagonia. Nine subfamilies are recorded from this region, including the monotypic subfamily Chilenomyiinae which is endemic for the area. Of the 111 known genera from South America, 53 are present in the study area. Approximately 20 % of the genera recorded from Patagonia are endemic to the region. One hundred and seventy seven species of Chironomidae have been recorded from Patagonia, 98% of them are endemic, 3 % are considered *nomina dubia* and about 17 % require revision since their systematic position is doubtful. Taking the Andean cordillera as a boundary, 55 species are endemic to the eastern side, 57 are endemic to the western side and 65 are common to both sides. The original descriptions of the chironomid species from Patagonia are based mainly on males (n=58) or males and females (n=43) which represents 57 % of the total. The rest of the species have been described using different combinations of stages from their life cycle (Donato et al., 2009). The Patagonian chironomid fauna shows affinities with the Australasian region (i.e. with a strict transantarctic distribution) exemplified by the Chironominae genera *Megacentron* Freeman and *Riethia* Kieffer, the Podonominae genus *Rheochlus* Brundin, and the Orthocladiinae genera *Austrocladius* Freeman, *Botryocladus* Cranston & Edwards, *Parapsectrocladius* Cranston, *Rhinocladius* Edwards and *Stictocladus* Edwards. Based on the material collected, mounted and studied so far during the last three years, we have found 16 new species that are waiting for more data (mostly concerning their immature stages) before they are described.

References

- Cabrera, A.L. & Willink, A. 1973. *Biogeografía de América Latina*. Monografía 13. Serie de Biología. Secretaría General de la Organización de los Estados Americanos. Washington DC. USA. 120 pp.
- Donato, M., Massaferrero, J. & Brooks, S.J. 2008. Chironomid (Chironomidae: Diptera) checklist from Nahuel Huapi National Park, Patagonia, Argentina. *Revista de la Sociedad Entomologica Argentina* 67: 163-170.

- Donato, M., Massaferrero, J. & Brooks, S.J. 2009. Estado del conocimiento taxonómico de la fauna de Chironomidae (Diptera: Nematocera) de la Patagonia [Current state of the taxonomic knowledge of the Chironomidae fauna (Diptera: Nematocera) of Patagonia]. *Revista de la Sociedad Entomologica Argentina* 68 (1-2): 187-192.
- Edwards, F.W. 1927. Insect collecting in the Southern Andes. *Natural History Magazine* 1: 111-125.
- Edwards, F.W. 1931. *Chironomidae. Diptera of Patagonia and South Chile, based mainly on material in the British Museum (Natural History)*. Part 2, Nematocera. London: Printed by Order of the Trustees. pp. 233- 331.
- Morrone, J.J. 2001. *Biogeografía de América Latina y el Caribe*. M&T-Manuales & Tesis SEA, vol. 3. Zaragoza, 148 pp.
- Olson, D.M., Dinerstein, E., Wikramanaya, E.D., Burgess, N.D., Powell, G.V.N., Underwood, E.C., D'Amico, J.A., Itoua, I., Strand, H.E., Morrison, J.C., Loucks, C.J., Allnutt, T.F., Ricketts, T.H., Kura, Y., Lamoreux, J.F., Wettengel, W.W., Hedao, P. & Kassem, K.R. 2001. Terrestrial Ecoregions of the World: A New Map of Life on Earth. *Biosciences* 51 (11): 933-938.
- Ringuelet, R.A. 1961. Rasgos fundamentales de la Zoogeografía de la Argentina. *Physis* 22 (63): 151-170.
- Sclater, P.L. 1858. On the general geographical distribution of the members of the class Aves. *Zoological Journal of the Linnean Society* 2: 130-145.
- Takhtajan, A. 1986. *Floristic regions of the world*. University of California. Berkeley, USA. 522 pp.
- Wallace, A.R. 1876. *The Geographical Distribution of Animals*. Macmillan, London.

DEVELOPMENT OF A NEW DESIGN OF AN INSECTARY MODEL FOR REARING AND ENVIRONMENTAL ASSESSMENT STUDIES ON CHIRONOMID MIDGES

B. B. Nath¹, Leena Thorat¹, Ananda Babrekar¹, Kailash Datkhile¹, Ashok Ghadge¹,
Rahul Gaikwad¹ & D. C. Meshram²

¹Centre for Advanced Studies, Department of Zoology, University of Pune, Pune-411007, India, bbnath@unipune.ernet.in, leenathorat@gmail.com, babrekar@gmail.com, kddatkhile@gmail.com, ghadgeas008@gmail.com & rahul0880@gmail.com

²Department of Geology, University of Pune, Pune-411007, India, dcmeshram@unipune.ernet.in

Chironomus ramosus Chaudhuri (1992), is one of the common tropical midges that form an abundant group of sediment-dwelling macro invertebrates of slow-flowing freshwater ecosystems of the Indian subcontinent, vulnerable to numerous fluctuating environmental conditions. Midges not only play an important role in maintaining a balanced aquatic ecosystem, being a major prey for the invertebrate and vertebrate fauna and preying upon lower invertebrates, but are also important rate determining conduits useful as indicators for water quality bioassessment (Merritt et al. 2009). In recent times, chironomid midges have emerged as a useful model system for environmental biomonitoring studies of acute and chronic

toxicity of large number of pollutants of the aquatic ecosystem.

Although laboratory based studies may provide a preliminary assessment of various biotic and abiotic factors circulating in the environment, they are mere auxiliary tools to the interpretation of *in situ* bioassays. For this reason, assessment of the responses of test organisms to laboratory exposures along with field based studies would help minimize the biases associated with lab-to-field extrapolation. Unfortunately, very few attempts have helped clarify this issue. A few of such studies have exploited midges, thus providing a better understanding of whether *in situ* and laboratory tests demonstrate comparable results (Faria et al. 2007). However, the absence

of an *in situ* control group of the biota is one of the limitations noticed in many contemporary experimental designs.

In an attempt at overcoming the shortcomings of comparative studies comprising laboratory and field samples, our laboratory team has come up with a novel *ex situ* technique suitable for rearing and carrying out simulation studies on *C. ramosus* in an insectary. This insectary is specially designed to meet the conditions as similar as possible to the prevailing natural environment. Further, rearing is carried out in small water tanks supplemented with phytoplankton and protists, thus mimicking the conditions in a lentic water body. Of the two tanks inside the insectary, one serves for *ex situ* control studies and the other for corresponding experimental studies in which stressor concentrations can be manipulated as per the requirement of the experimental design. The insectary is kept covered with a shade net in order to prevent the entry of other insects. The design of the insectary is depicted in Fig.1 a-c. This technique has been successfully standardized for the maintenance and breeding of *C. ramosus* and is cost effective involving less labour input. To

the best of our knowledge, there are no reports of similar techniques practised for chironomid midges where natural populations can be maintained and subjected to natural seasonal variations.

The tanks were constructed of clay bricks reinforced with concrete columns. To begin with, egg masses collected from the field were reared and developmental stages were validated for taxonomic identification using morphological and cytotoxic keys (Chaudhuri et al. 1992, Nath and Godbole 1997). To initiate rearing, ≈ 100 late fourth instar larvae each were transferred to rearing tubs ($\text{Ø}=35$ cm) placed in a net cage under laboratory conditions (Fig. 1 d), which were maintained exactly as described by Nath and Godbole (1998). Similarly, ≈ 100 larvae were transferred to the insectary tanks, covered with net cages and the base of the tanks was layered (2 cm) with silt and finely ground moss that served as soft substratum for tube building. Special food formulation (Nath and Godbole 1998) was supplied to the developing cultures every two days.




Fig. 1: a) Blue print of the insectary showing the location of the rearing tanks within the insectary. b) Interior view of the insectary showing the two rearing tanks. The insectary is shown covered with a shade net. c) Interior view of the insectary showing the two rearing tanks covered with net cages and each filled with about 130 litres of water. d) Special laboratory facility for rearing populations under ambient laboratory conditions.

The absence of predators of *C. ramosus*, placed it at the last trophic level of the food chain in the rearing tanks. It was found unnecessary to change the water of the tanks on a routine basis, thus saving both time and effort. However, whenever cleaning of the ponds was demanded, the water was flushed out through a small outlet present at one of the basal corners of the tanks. The desired level of water in the tanks was maintained through appropriate replenishments.

Our observations indicated that the life cycle of *C. ramosus* from egg to adult in the laboratory reared populations was completed in about 36 days, while it took about 33 days for the insectary reared population. The differences of growth parameters found in the two groups was found to be negligible (ANOVA, $P > 0.05$) (Table1).

Table 1: Comparison between lab reared (LR) and insectary reared (IR) populations for various growth, developmental and emergence parameters. d = duration in days.

Parameter	LR population	IR population
Hatching of eggs	2d	2d
1 st to 4 th instar stage	29 ± 1.25 d	25 ± 2.06 d
Pupal stage	3 ± 0.11d	3 ± 0.21 d
Adult life span	2.5 ± 0.68 d	3 ± 0.32 d
# of swarming adults on the day of egg laying	20 ± 3.25	25 ± 2.66
Sex ratio on the day of egg laying	1:1	1:1
Total egg masses obtained	6	9

The suitability of this novel technique proved successful when egg masses (i.e. 1st generation) were obtained from the founder population. When reared in a similar manner, these egg masses perpetuated to successive generations (so far, 4 generations have been successfully reared in the tanks). Therefore, the rearing conditions maintained in the insectary showed future potential for undertaking genetic toxicological studies. Most of the ecotoxicological research has been aimed at investigating the responses of field samples under laboratory experimental regimes. But what these studies lack is an *in situ* understanding and implications of the stressors. The novel approach adopted by us, would provide

an alternative design for *ex situ* assessment of toxicants using chironomid midges as model systems and gaining useful insights to queries of the ecological implications of the abiotic factors that prevail in the natural environment and whose role in the various developmental processes cannot be played-out. We hope that the model of the insectary presented here, would attract the attention of chironomidologists for designing simulation experiments, often required in environmental assessment studies.

Acknowledgements

BBN and DCM acknowledge the financial support received from ISRO-UoP research project (# 80) and the infrastructural facilities from the Department of Zoology, University of Pune. The authors also acknowledge the technical suggestions received from Mr. Yeshwant B. Kulkarni.

References

- Chaudhuri, P.K., Das, S.K. and Sublette, J.E. 1992. Indian species of genus *Chironomus* Miegen (Diptera: Chironomidae). - *Zoologische Jahrbücher für Systematik* 119: 1 - 51.
- Faria, M.S., António J.A., Nogueira, and Amadeu M.V.M. Soares. 2007. The use of *Chironomus riparius* larvae to assess effects of pesticides from rice fields in adjacent freshwater ecosystems. - *Ecotoxicology and Environmental Safety* 67: 218-226
- Merritt, R.W., Courtney, G. W. and Keiper, J. B. 2009. Diptera. In: Resh V. H. and Carde C.R. (Eds.). *The Encyclopedia of Insects*. Elsevier, Amsterdam, pp 284- 299.
- Nath B.B. and Godbole N.N. 1997. Chromosomal characterization of tropical midge. - *Cytobios* 91: 25-31.
- Nath B.B. and Godbole N.N. 1998. Technique for mass rearing of Indian *Chironomus* species. - *Studia Dipterologica* 5: 187-193.

CURRENT RESEARCH

SYNONYMY OF *CHIRONOMUS PLUMATISSETIGERUS* TOKUNAGA, 1964, WITH *CHIRONOMUS CIRCUMDATUS* KIEFFER, 1916

Jon Martin¹ and Sumitra Saxena²

¹Department of Genetics, University of Melbourne VIC 3010, Australia, j.martin@unimelb.edu.au
²S.G.T.B. Khalsa College, University of Delhi, Delhi, India

Recently, Pramual *et al.* (2008) noted the synonymy of *Chironomus plumatisetigerus* with *Chironomus circumdatus*, based on the information of the web page of Martin (2007). That web page is not a recognised publication and, further, the synonymy is not indicated in either the title or abstract of the paper. This note is therefore intended to provide a more formal notification and basis for the synonymy.

C. circumdatus was described by Kieffer (1916) from Formosa (Taiwan), and the type is presumably lost. However, the species has been redescribed by a number of people, of whom the most relevant here is that of Chaudhuri *et al.* (1992), since material used for cytological studies in India (Kumar and Gupta 1990; Tripathi *et al.* 2004) is based on this description.

C. plumatisetigerus was described by Tokunaga (1945) from several islands of Micronesia. Material identified as this species was found in Australia (Bugledich *et al.* 1999), and this material was the basis for publications on the cytology of this species in Thailand (Kuvangkadilok 1985) and India (Saxena 1995).

Comparison of the published chromosome maps from the studies under these two names clearly showed that the banding patterns and most of the inversion polymorphisms were identical, indicating that all studies were of the same species. This cytologically defined species becomes *C. circumdatus*, since that name has priority, and so *C. plumatisetigerus* must be considered as a junior synonym.

References

Bugledich, E.-M.A., Cranston, P.S. and Martin, J. 1999. Diptera: Nematocera: Chironomidae. *In* Wells, A. & Houston, W.W.K. (Eds.)

Zoological Catalogue of Australia, Vol. 30.1, CSIRO Publishing, Melbourne, pp. 112-158.

Kieffer, J.J. 1916. Tendipedides (Chironomides) de Formose conservés au Muséum National Hongrois de Budapest et déterminés par J. J. KIEFFER. - *Annales historico-naturales Musei nationalis Hungarici* 14 : 81-121.

Kumar, A. and Gupta, J.P. 1990. Cytogenetic studies of *Chironomus circumdatus* from India (Diptera: Chironomidae). - *Genetica* 82: 157-163.

Kuvangkadilok, C. 1985. Cytogenetic studies of *Chironomus plumatisetigerus* (Diptera: Chironomidae) in Thailand. - *Journal of the Sciences Society of Thailand* 11: 37-45.

Martin, J. 2007. Australian *Chironomus* species. Available from <http://www.genetics.unimelb.edu.au/Martin/AustChironfile/AustChiron.htm> (accessed 14 July 2009)

Pramual, P., Gomontean, B., Buasay, V., Srikhamwiang, N., Suebkar, P., Niamlek, C., Donsinphoem, Y. and Chalath-Chieo, K. 2008. Population cytogenetics of *Chironomus circum-datus* Kieffer, 1921 (Diptera, Chironomidae) from Thailand. - *Genetica* 135: 51-57.

Tokunaga, M. 1964. Diptera, Chironomidae. - *Insects of Micronesia* 12: 485-628.

Tripathi, N.K., Sharma, O.P. and Khanna, P. 2004. Studies on the polytene chromosomes of *Chironomus circumdatus* (Diptera: Chironomidae) from Jammu region. - *Proceedings of the National Academy of Sciences India, Section B, Biological Sciences* 74: 1-6.

A BRIEF ANALYSIS OF MORPHOLOGY AND TERMINOLOGY FOR THE GENUS *ABLABESMYIA* (TANYPODINAE)

Analía C. Paggi¹, Augusto Siri¹ and Caroline Silva Neubern de Oliveira²

¹ Instituto de Limnología "Dr. R.A. Ringuelet" (CONICET, UNLP) C.C. 712, 1900 La Plata, Buenos Aires, Argentina. E-mail: anpaggi@ilpla.edu.ar and augusto@ilpla.edu.ar

² Departamento de Zoologia, Universidade Federal do Paraná. Caixa Postal. 19020, Curitiba, Paraná, Brasil. E-mail: cneubern@yahoo.com.br

The genus *Ablabesmyia* is well represented in both the north and south Americas, though the latter region is inadequately studied. A few months ago, the authors of this note made contact to exchange type material and literature on the known species of *Ablabesmyia* from South America. Almost immediately, we faced the problem that we were referring to the same structures by different names, so we decided to sort and organize the different terminologies used along with the most important bibliography of this genus. Furthermore, we wanted to share the misunderstandings we faced with others who want to study *Ablabesmyia* so they can learn from our mistakes.

Megaseta of the adult male gonostylus

When describing the male hypopygium of *Ablabesmyia*, Johannsen (1946: 271) wrote the following: "styles of the hypopygium about as those of a typical *monilis* except that the articulated pre-apical stylet is straight and acute at apex". Roback some years later (1959: 116) separated the main subgenera, *Ablabesmyia s. str.* and *Karelia*, by the characteristic "preapical stylet pointed (P) or expanded (E)", however, the same

author used different names for the same structure later in the same paper: "preapical spine of the distyle" and "preapical spur of the distyle has retained the simple apex" (p. 117). Roback (1971: 356, 357 and 365) employed the name "preapical spur", and in 1982 (p. 106) "spur of dististyle". In 1983, the same author used "gonostylus spur expanded or sharp" (Roback 1983: 237, table 1).

Later, Roback (1985) introduced the term "megaseta", in his subgeneric adult diagnosis of the subgenera *Karelia* (p. 169) and *Ablabesmyia* (p. 178), probably adopting the nomenclature of Sæther & Sublette (1983) for the homologous structure in *Orthocladinae*. Unfortunately, Murray & Fittkau (1989), in their *Tanypodinae* diagnosis for the Holarctic genera, erroneously used the term "megaseta" to refer to the apex of the gonostylus: "gonostylus simple...distally with megaseta, mostly without but occasionally with simple or apically expanded subterminal seta (*Ablabesmyia*)..." (p. 38); followed by "gonostylus as long as or slightly longer than gonocoxite, slender and tapering; megaseta cochleariform with distally expanded or pointed subterminal seta" (p.42). This misunderstanding confused us and generated doubts on which structure actually is the megaseta.




Figure 1. Distal portion of gonostylus. a) extracted from Paggi & Añón Suarez 2000. b) extracted from Paggi 1987

In order to clarify and unify the use of terms describing the subterminal seta on the gonostylus of Tanypodinae, we suggest that the name “megaseta” (Roback 1985) should be used. The megaseta lies parallel to the “hooded or cochleariform” apex of gonostylus (Fig. 1). In descriptions of *Ablabesmyia hypopygia*, it is very important to consider the shape of the megaseta (expanded or sharpened) to differentiate the subgenera and to avoid the wrong interpretation. Furthermore, it is very important to place the structures in their correct position during slide mounting, keeping the apex of the gonostylus and megaseta a little separated from another. The gonostylus should preferably be mounted in lateral view to avoid possible misinterpretations (Fig. 1).

The Aedeagal complex

With regard to the aedeagal complex we also recorded different morphology terminology for the same structures. Roback (1959, 1971) described this complex as being formed of four pairs of structures (aedeagal blades, dorsal lobes, lateral filaments and lateral lobes). Later, Sæther (1980) named these structures volsellas (superior, median and inferior volsellas), but he did not mention the “lateral lobes” described by Roback.

It is possible that they were hidden by the other structures or absent in the examined specimens. Roback (1983) did not agree with Sæther (1980) on the homology of the lobes with the volsellas of the other subfamilies. He introduced the term basidorsal lobe of the gonocoxite and suggested that it might be the true inferior volsella of the other subfamilies. We therefore think it is appropriate to describe the aedeagal complex, to accurately name the different structures, and choose one set of terminology to avoid confusion. This has been attempted in Fig. 2.

Maxillary palps in adults and larvae

Older works on chironomids, describe four palpomeres in the maxillary palp of the adult male. Recent papers describe the palps to consist of 5 segments, following the works by Sæther (1971, 1980). The new interpretation acknowledges the first palpomere as a small, basal, weakly sclerotized segment that is deprived of setae. This means that what was previously considered to be the third palpomere, now is the fourth, and that the fourth palpomere (not the third) is shorter than the third (not the second as described by Fittkau 1962: 416 and Roback 1971: 354).




Figure 2. Aedeagal complex. a) *Ablabesmyia bianulata*, modified from Paggi 1987. b) *Ablabesmyia simpsonii*, modified from Roback 1985

Numbering of the larval palpal segments has likewise been done differently by authors and thus caused inconsistency in the calculation of the length ratio between the first and second palpal segments. Roback (1985, fig. 42) started the numbering from the apical palpomere while Epler (2001: 4.20) named the most basal palpomere segment 1. Since the number of palpomeres in *Ablabesmyia* larvae increase by the addition of segments basal to the ring organ, we suggest following Roback's (1985) numbering for the future. Thus the palpomere distal to the ring organ should be named P1.

Different use of morphology terminology can easily result in erroneous interpretation of identification keys, both at the subgenus and species level in *Ablabesmyia*. We hope that the above clarification of names for structures observed in this genus will aid in future nomenclature consistency.

Acknowledgments

The authors thank Bohdan Bilyj and Torbjørn Ekrem for their comments and suggestions on the manuscript.

References

- Epler, J. H. 2001. Identification Manual for the larval Chironomidae (Diptera) of North and South Carolina. A guide to the taxonomy of the midges of the southeastern United States, including Florida. Special Publication SJ2001-SP13. North Carolina Department of Environment and Natural Resources, Raleigh, NC, and St. Johns River Water Management District, Palatka, FL. 526 pp. <http://home.earthlink.net/~johneppler/index.html>
- Fittkau, E.J. 1962. Die Tanypodinae (Diptera, Chironomidae). Die Tribus Anatopyniini, Macropelopiini und Pentaneurini. - *Abh. Larvalsyst. Lnsekten*. 6: 1-435.
- Johannsen, O.A. 1946. Revision of the North American species of the genus *Pentaneura* (Tendipedidae: Chironomidae, Diptera). - *J. New York Entomol. Soc.* LIV. 4: 267-289.
- Murray, D.A. and Fittkau, E. J. 1989. The adult males of Tanypodinae (Diptera: Chironomidae) of the Holarctic Region – Keys and Diagnosis. In: Wiederholm, T. (Ed.), Chironomidae of the Holarctic region – Keys and diagnoses. Part 3. Adult males. - *Ent. Scand. Suppl.* 34: 37-123.
- Roback, S.S. 1959. The subgenus *Ablabesmyia* of *Pentaneura* (Diptera, Tendipedidae, Pelopiinae). - *Trans. Amer. Ent. Soc. Philad.* 85: 113-135.
- Roback, S.S. 1971. The adults of the subfamily Tanypodinae (= Pelopiinae) in North America (Diptera: Chironomidae). - *Acad. Nat. Sci. Philad.* 17: 1-410.
- Roback, S.S. 1982. The Tanypodinae (Diptera: Chironomidae) of Australia II. - *Proc. Acad. Nat. Sci. Philad.* 134: 80-112.
- Roback, S.S. 1983. *Ablabesmyia* (*Sartaia*) *metica* a new subgenus and species (Diptera: Chironomidae: Tanypodinae). - *Proc. Acad. Nat. Sci. Philad.* 135: 236-240.
- Roback, S.S. 1985. The immature Chironomids of the eastern United States.VI. Pentaneurini-genus *Ablabesmyia*. - *Proc. Acad. Nat. Sci. Philad.* 137, 2: 153-212.
- Sæther, O.A. 1971. Notes on general morphology and terminology of the Chironomidae (Diptera). - *Can. Ent.* 103: 1237–1260.
- Sæther, O.A. 1980. Glossary of chironomid morphology terminology (Diptera: Chironomidae). - *Ent. scand. Suppl.* 14: 1-51.
- Sæther, O.A. and Sublette, J. E. 1983. A review of the genera *Doithrix* n. gen., *Georthocladius* Strenzke, *Parachaetocladius* Wülker, and *Pseudorthocladius* Goetghebuer (Diptera: Chironomidae, Orthocladiinae). - *Ent. scand. Suppl.* 20: 1-100.

PALEOLIMNOLOGICAL EVIDENCE CONFIRMS THAT *PAROCHLUS STEINENII* (GERKE) IS NOT A RECENT INTRODUCTION TO THE ANTARCTIC PENINSULA REGION

Janelle T. Agius¹, John A.E. Gibson^{1,2*}, Eugenio Rico³ and Antonio Quesada⁴

¹*Institute for Antarctic and Southern Ocean Studies, University of Tasmania, Private Bag 77, Hobart, Tasmania, Australia. Email: John.Gibson@utas.edu.au*

²*Marine Research Laboratories, Tasmanian Aquaculture and Fisheries Institute, University of Tasmania, Private Bag 49, Hobart, Tasmania, Australia*

³*Departamento de Ecología, Universidad Autónoma de Madrid, 28049 Madrid, Spain*

⁴*Departamento de Biología, Universidad Autónoma de Madrid, 28049 Madrid, Spain*

* Corresponding author

Introduction

Chironomid remains are common in lake sediments, and have been used widely as indicators of change in climatic conditions. Far fewer studies, however, have made use of the palaeobiogeographical potential of these remains to investigate past species distributions and the origins of isolated faunas.

Two models have been suggested for the origin of Antarctic populations of the chironomid *Parochlus steinenii* (Gerke) (Allegrucci et al. 2005). It was originally suggested that this species could have been introduced to the Antarctic Peninsula region from farther north by human activities, beginning with whaling in the mid-to-late 1800s (Convey and Block 1996). This suggestion was predicated on the quite recent initial observation (1950s) of this species in the South Shetland Islands and the Antarctic Peninsula (Torres 1956).

Molecular studies indicated a different story. Analysis of 28S ribosomal RNA from three distinct populations of *Parochlus steinenii* suggested that the populations on the Antarctic Peninsula and the nearby South Shetland Islands had been separated from the other populations on a time scale of million of years, and that each population was limited to particular tectonic plates (Allegrucci et al. 2005).

In this short note we use palaeolimnological techniques to test these models, and conclude an ancient origin for the Antarctic Peninsula/South Shetland Island population of *Parochlus steinenii*.

Methods

The study site was a small lake on Byers Peninsula, Livingston Island (62°40'S, 61°00'W, Figure 1), that has been the centre of a concerted scientific study (Toro et al. 2007), and which has become known as Limnopolar Lake. See www.aslo.org/photopost/showphoto.php/photo/706/ppuser/158 for a picture of the lake. Livingston Island is in an active volcanic area, with nearby Deception Island having erupted as recently as 1969 (Pallàs et al. 2001).

During the Antarctic field season of 2002/2003 a 29.2 cm sediment core was obtained from Limnopolar Lake using a modified hybrid Glew corer and Kajak corer. The core did not reach bedrock. The core was sectioned on site into 0.2 cm sections (0-10 cm), 0.5 cm sections (10-25 cm), or 1.0 cm sections (25-29 cm). The sections were then transferred into labelled whirlpak bags, and stored at 4°C until analysed.

Animal remains were isolated by placing approximately 1 g of wet sediment in a 20 ml scintillation vial and adding distilled water to disperse the sediment. The sediment was sieved on 44 µm and 100 µm mesh, and the material retained on the sieves was treated with Rose Bengal to stain biological components. The material was examined under a dissecting microscope and chironomid remains isolated for further study.

Radiocarbon dating of the sediments, estimation of the relative abundance of aquatic moss (on an arbitrary scale), and measurement of abundance of tephra (microscopic shards of volcanic glass) were undertaken as described elsewhere (Agius 2006).




Figure 1. Map of southern South America and the Antarctic Peninsula showing the location of Livingston Island, as well as other areas inhabited by *Parochlus steinenii* (Terra del Fuego and continental South America, South Georgia, South Shetland Islands, Antarctic Peninsula mainland).

Results

Chironomid remains were encountered at abundances up to 180 per g dry weight (g_{dw}^{-1}). The remains, which were typically of the order of 200 μm in length, were mainly portions of adults and discarded pupae, with many anatomical




Figure 2. Photographs of representative *Parochlus steinenii* remains: (a) head capsule and eyes of adult; (b) wings and legs of pupa; (c) tail of pupa; and (d) adult thorax.

features present (Figure 2). The remains could be identified by comparison to literature figures, in particular the figure of adult and pupa in Wirth and Gressitt (1967). Identification was confirmed by P. Convey, British Antarctic Survey, UK. Somewhat surprisingly, no larval head capsules were observed.

The distribution within the core (Figure 3) exhibited peaks at depths of 8 cm, 10-15 cm and 25 cm separated by periods of low abundance. The distribution of the chironomid remains showed qualitatively similar trends to that of aquatic moss, with both the moss and the chironomids less abundant after periods of significant tephral input.

The age of the sediments was difficult to determine due to contamination with old carbon, possibly through partial melting of permafrost (J. Gibson and A. Quesada, unpublished results). The most parsimonious interpretation of the data suggested that the core covered approximately 2000 years of sedimentation, but this figure must be viewed with caution.




Figure 3. Abundance of *Parochlus steinenii* remains (per gram dry weight) in a sediment core from Linnopolar Lake as a function of depth. Also shown are the abundance of tephra shards and a qualitative estimate of the amount of dead moss present in the sediment. The time span covered by the sediment core is probably of the order of 2000 years.

Discussion

The occurrence of remains of *Parochlus steinenii* at nearly all depths sampled within the sediment core indicates that the species was present throughout the time the sediment was deposited. While the precise time interval the core represents cannot be determined, it is certain that the chironomid has been present on Byers Peninsula for an extended period, and that the suggestion that the species was introduced by whalers can be almost certainly refuted. The observed distribution within the sediment is consistent with the conclusions of Allegrucci et al. (2005), who suggested an ancient origin for this species and therefore the presence of glacial refugia either on the Antarctic Peninsula or within the South Shetland Islands. Recognition of similar glacial refugia across Antarctica is increasing (Convey et al. 2008)

The distribution of the remains in the core gives some indication of the habitat preferences of the species: it appears to be more abundant when significant moss growth occurs in the lake. Toro et al. (2007) reported that *Parochlus steinenii* was most likely to be present in Livingston Island

lakes that also contained benthic mosses, which is consistent with this conclusion. Abundance appears to be negatively impacted by deposition of tephra from nearby volcanic eruptions, which would smother food sources. Similar observations have been made during the study of a Chilean lake impacted by tephral deposition (Urrutia et al. 2007). However, it may also be that these apparent correlations are artefacts stemming from changes in sedimentation rates as a result of tephra and other volcanic inputs. The absence of larval head capsules in the sediment is surprising, as larvae inhabit Linnopolar Lake (Toro et al. 2007). The conclusion to be drawn from this observation is that the head capsules of *Parochlus steinenii* are not preserved in the sediment.

Acknowledgements

We thank M. Toro for collecting the core and kindly providing the sediments for this research, P. Convey for confirming the identity of the chironomid remains, and the Australian Institute for Nuclear Science and Engineering for funding the attempts at dating the sediment core (Grants 05063 and 07066). The Ministerio de Educación y Ciencia (Spain) also funded parts of this work

with the grants REN2000-0435-ANT and CGL2005-06549-CO2-01.

References

- Agius, J.T. 2006. Tephra and ecological studies of Limnopolar Lake, Byers Peninsula, Livingston Island. Honours thesis, University of Tasmania. Available from: <http://eprints.utas.edu.au/8815/>
- Allegrucci, G., Carchini, G., Todisco, V., Convey, P. and Sbordoni, V. 2005. A molecular phylogeny of Antarctic Chironomidae and its implications for biogeographical history. - *Polar Biology* 29: 320-326.
- Convey, P. and Block, W. 1996. Antarctic Diptera: ecology, physiology and distribution. - *European Journal of Entomology* 93: 1-13.
- Convey, P., Gibson, J. A. E., Hillenbrand, C., Hodgson, D. A., Pugh, P. J. A., Smellie, J. L. and Stevens, M. I. 2008. Antarctic terrestrial life – challenging the history of the frozen continent? - *Biological Reviews* 83: 103-114.
- Pallàs, R., Smellie, J.L., Casas, J.M. and Calvet, J. 2001. Using tephrochronology to date temperate ice: correlation between ice tephrae on Livingston Island and eruptive units on Deception Island volcano (South Shetland Islands, Antarctica). - *The Holocene* 11: 149-160.
- Torres, B.A. 1956. Primer hallazgo de Tendipedidos alados en lae region Antartica. Podonominae, una nueva subfamilia para la citada region. - *Anales de la Sociedad Cientifica Argentina* 161: 41-52.
- Toro, M., Camacho, A., Rochera, C., Rico, E., Banon, M., Fernandez-Valiente, E., Marco, E., Justel, A., Avendano, M.C., Ariosa, Y., Vincent, W.F., and Quesada, A. 2007. Limnological characteristics of the freshwater ecosystems of Byers Peninsula, Livingston Island, in maritime Antarctica. - *Polar Biology* 30: 635-649.
- Wirth, W.W. and Gressitt, J.L. 1967. Diptera: Chironomidae (Midges). - *Antarctic Research Series* 10: 197-203.
- Urrutia, R., Araneda, A., Cruces, F., Torres, L., Chirinos, L., Treutler, H.C., Fagel, N., Bertrand, S., Alvial, I., Barra, R., and Chapron, E. 2007. Changes in diatom, pollen, and chironomid assemblages in response to a recent volcanic event in Lake Galletue (Chilean Andes). - *Limnologica* 37: 49-62.
-

SHORT COMMUNICATIONS

Chironomidae photo database of the Sasa collection opened on the home page of the National Museum of Nature and Science, Tokyo

Tadashi Kobayashi

Ex-senior researcher of Institute for Environmental and Social welfare Studies, Kurobe

E-mail: tadkoba@k.email.ne.jp

As presented in Chironomus Newsletter No.19 (2006) by Dr. Toshio Iwakuma, the late Professor Dr. Manabu Sasa (1916-2006) began his studies on Chironomidae in 1974, when he was 58 years old, while vice director of the National Institute for Environmental Studies (NIES). In those days, nuisance chironomids were causing a big problem around Kasumigaura Lake, which is located near the Institute. When he began his studies, he visited Dr. M. Tokunaga, still alive at Kyoto, and was given charge of Dr. Tokunaga's library.

Dr. Sasa described 1,151 species of Chironomidae as new. Information on the types including the current names (as of November, 2007) are listed in the Appendix of YUSURIKA Supplement 2 ('In Memory of Professor Dr. Manabu Sasa') delivered by the Japanese Group for Chironomidae Studies in November, 2007. If anyone wants to see the Appendix, the author could send it to him as a pdf file.

After Dr. Sasa passed away in April, 2006 at Kurobe, Toyama, the part of his library relevant to the Chironomidae and about fifty thousand slide specimens were entrusted to the author. His library of 1,105 items was delivered to Dr. K. Hirabayashi of Shinshu University with the list in July, 2006. All slide mounted specimens of Chironomidae used by Dr. Sasa had been transferred to NIES from Kurobe in September, 2000. After this, in February, 2003 they were transferred to the annex of the National Museum of Nature and Science (NMNS), Shinjuku, Tokyo, as The Sasa Collection. The author had arranged the collection and compiled the database. In autumn, 2004, he was asked to take photos of all the types by Dr. A. Shinohara of the Museum. The photos consisted of the type slides themselves, and microscope photos of the male hypopygium, wing, and so on.

National Museum of Nature and Science
TYPE SPECIMEN DATABASE

Type Specimen Search: [Animals] > [Diptera] Back Top

Paratanytarsus kuramacircus M. Sasa, 1989 Chironomidae

Specimens Details

Specimen No.	165:071
Type Status	Holotype
Locality	Kurama River, Kyoto Pref., Japan
Collected Date	11 Oct., 1988
Collector	M. Sasa
Sex	
Remarks1	

Image

	hypopygium	hypopygium	thorax
	[Large]	[Large]	[Large]

Bibliography

Status	Original description
Title	Annex. Chironomid midges of some rivers in western Japan.
Author(s) of Paper	
Publication	Research Report of Toyama Prefectural Environmental Pollution Research Center, 45-110.
Publication Date	1989
Description on Page(s)	p. 50-51, fig. 26
Remarks2	4666

Screen shot of the Type Specimen Database

The photo database opened recently on the homepage of NMNS <http://www.type.kahaku.go.jp/TypeDB/> is based on these photos. If it is absolutely impossible to see the needed characters in the available photographs, a request can be made to Dr. A. Shinohara shinohar@kahaku.go.jp for loan of the types.

Next International Chironomidae Symposium to be held in Trondheim, Norway

The 18th International Symposium on Chironomidae Research will be hosted by the Museum of Natural History and Archaeology at the Norwegian University of Science and Technology in Trondheim, Norway in **2011**. We are very pleased that our bid to hold the next chironomid meeting was so well received by the recent symposium in Tianjin, and will do our best to make this a memorable event both scientifically and socially.



The city of Trondheim as seen from the city forest

Trondheim is situated in middle Norway (~ 63° N) and is Norway's third largest city with almost 170 000 inhabitants. It is home to one of the largest universities (approximately 18 000 students) and the oldest natural history collection in the country which can be dated back to the work of bishop Gunnerus (1718-1773). [The Museum of Natural History and Archaeology](#) was founded as the [Royal Norwegian Society of Sciences and Letters](#) as early as 1760.

Nature research and management is strong in Trondheim, and institutions like the University, The Norwegian Institute of Nature Research, the Directorate of Nature Management and The Norwegian Biodiversity Information Centre contributes to a considerable focus on biodiversity in our region.

Like most Norwegian cities, the city centre of Trondheim is close to nature. We hope that our meeting in Trondheim will give you the opportunity to experience some regional culture and wild life (don't forget your net) and look forward to see you here. For more information on the city and its surroundings, please see www.trondheim.no and www.visit-trondheim.com.

Torbjørn Ekrem, Elisabeth Stur & Kaare Aagaard

A World Catalogue of Chironomidae (Diptera). Part 1. Buchonomyiinae, Chilenomyiinae, Podonominae, Aphroteniinae, Tanypodinae, Usambaromyiinae, Diamesinae, Prodiamesinae and Telmatogetoninae

Patrick Ashe¹ & James P. O'Connor²

¹33 Shelton Drive, Terenure, Dublin 12, Ireland, patrick.ashe@upcmail.ie

²National Museum of Ireland, Kildare Street, Dublin 2, Ireland

Part 1 of A World Catalogue of Chironomidae (Diptera) is expected to be published in December 2009. A quote from the publisher has now been received and the price for each copy has now been fixed at Euro 42 to which the relevant amount of postage must be added. The number of pages will be approximately 370-400 and the volume will weigh about 1.3 to 1.4 kg. It is possible therefore to estimate the postage cost (Airmail only) to anywhere in the world. The total cost of a copy plus postage and packing can be determined from the table given below.

	Ireland (incl. Northern Ireland)	Britain	Europe	Rest of the World
Postage	Euro 8.00	Euro 13.00	Euro 13.00	Euro 18.00
Catalogue part 1	Euro 42.00	Euro 42.00	Euro 42.00	Euro 42.00
Total	Euro 50.00	Euro 55.00	Euro 55.00	Euro 60.00

The Catalogue is being published by The Irish Biogeographical Society in association with The National Museum of Ireland.

Once the Catalogue is published and available for sale it will be announced on the website of The Irish Biogeographical Society: <http://www.irishbiogeographicalsociety.com/>

The Bank which handles The Irish Biogeographical Society account will be consulted to determine the easiest and cheapest method for receiving payment from abroad – this information will be included on the website when the Catalogue is published. Money received from the sale of Part 1 will be used to help pay some of the cost of Part 2.


Anyone wishing to order a copy can contact the senior author, Dr Patrick Ashe, by e-mail: patrick.ashe@upcmail.ie

New books

Chironomidae larvae – Biology and Ecology of the Chironomini

H.K.M. Moller Pillot is one of those leading chironomidologists in the Netherlands, if not the leading person, who have a long experience in taxonomy and ecology of chironomid larvae. His splendid identification keys have been published in the early 1980s, but exclusively in Dutch. Those keys were the basis of the development of chironomid research in the Netherlands. With the new series “*Chironomidae Larvae of the Netherlands and Adjacent Lowlands*”, published in English, his knowledge now becomes accessible for the international scientific world. The first book of the series, “*Chironomidae larvae - General ecology and Tanypodinae*”, written by H.J. Vallenduuk and H.K.M. Moller Pillot, was published in 2007. It was quite original as it includes, together with an ecological review, a lot of very new, unknown and practical characters making the identification of Tanypodid larvae much easier, sometimes already possible by the stereomicroscope alone (see review by Elisabeth Stur in *Chironomus newsletter of Chironomidae research* volume 20).

The bulk of the second book in the series, “*Chironomidae larvae – Biology and ecology of the Chironomini*”, is mainly limited to larval autecology (Pseudochironomini are also included, i.e. *Pseudochironomus prasinatus*). It is a fundamental synthesis per species(-group) in a rather constant pattern of topics (as far as data are available): life cycle, microhabitat, feeding, swarming and oviposition, dispersal, densities, water type (current, dimensions, permanence, shade, ...), pH, salinity, trophic conditions and saprobity, etc. When other specific environmental requirements are known, they are treated separately. Distribution in Europe is mainly based on Sæther & Spies (2004) to which specific data concerning the Netherlands are added. The book is limited to Dutch species; concerning their ecology, the situation in other countries is also taken into account, but the accent still remains on Central European lowland. The accounts are concise and they also include a lot of hidden literature and unpublished data; the work of Russian chironomidologists, e.g., is largely considered.



The last chapter (p. 239-247) concerns four tables with biological and ecological properties of Chironomini in the Dutch situation: (1) general ecology of each species(-group): number of generations, flying period, duration of larval development, diapausing instar stage, number of eggs, food and substrate of the larvae, (2) occurrence in relation to saprobity and oxygen concentration, (3) in relation to pH and chlorinity, and (4) in relation to water velocity and permanence. Each variable in table 2-4 is subdivided into several classes (mostly five) and a score from 0 to 10 is given to the relative occurrence of each species (-group) in these classes. It is not clear how these occurrence scores have been calculated and what confidence may be given to them: the author must have had a lot of (unpublished) data at his disposal, both from himself and from his network, but the exact number for each species is not given.

“*Chironomidae larvae – Biology and ecology of the Chironomini*” does not contain identification keys, but warnings concerning misidentifications and indications of correct diagnostic descriptions in literature are mentioned; a few drawings sometimes illustrate specific characters. In any case, these remarks are so welcome that the reader becomes anxious to have the revised keys already at his disposal; the publication of these keys is foreseen for the next volume in this series.

One may regret that the book (and series) is mainly limited to species of the Netherlands and that even some aspects of their ecology only refer to Western European lowland conditions. Including data from other countries in the ecological tables of chapter 4 was not an option: (1) in different climatic conditions the species may react in another way, e.g. living in cool lakes in northern regions versus running waters in temperate regions, and (2) a dominance of Dutch data compared to other countries could be expected, which would distort our view on the ecology of the species at e.g. a European level. Moreover, the author is mainly involved in alpha-taxonomy, faunistics and environmental requirements of chironomid larvae in

the Netherlands, the species with which he has been involved on a day-to-day basis. An extended edition including other European regions would have postponed, eventually cancelled this publication.

These short words to stress the high value of this publication. It is a must for everybody involved with chironomids. The “high” price compared to the volume of this book may not be taken into consideration: the content of this book is important and a gain of time for all of us who work with European chironomids.

Dr. Boudewijn Goddeeris, K.B.I.N., Vautierstraat 29, 1000 Brussels, Belgium.

Book reference

Moller-Pillot, H.K.M. 2009. *Chironomidae larvae II – Biology and Ecology of the Chironomini*, KNNV Publishing Zeist, The Netherlands, pp. 270. ISBN 978 90 5011 3038

How to order

Order from: KNNV Publishing, PO Box 310, 3700 AH Zeist, The Netherlands. Fax: +31 30 2368907.
www.knnvuitgeverij.nl Email: info@knnvuitgeverij.nl.

Price: 69,50 € postage not included.

CURRENT BIBLIOGRAPHY: 06 OCT. 2008 - 20 OCT. 2009

Odwin Hoffrichter

Institut für Biologie I, Albert-Ludwigs-Universität Freiburg, Hauptstr. 1, D-79104 Freiburg, Germany
Email: odwin.hoffrichter@biologie.uni-freiburg.de

In addition to supplements to the last two preceding years the publications of the present year are shown: almost one citation per day, which gives the average speed of publishing with regard to chironomids. The compilation was achieved, as usual, from many sources: databases, tables of contents of journals, references and citations of papers, inspection of many periodicals, lists and pdf's provided by authors (thanks to you!). In particular, publisher issued search alerts proved to be rich in results. As a rule, only printed titles are reported here with the occasional, but obviously increasing, exception of online-only journals (PLoS or BioMed Central journals, e.g.). Titles announced "in press", even with available DOI numbers, are not considered before printing. In general, online information should be retrieved elsewhere; best check the chironomid home page for eventual references regularly, or use individual websites with a host of chironomid-related data. Publications using chironomids as prey or food for animals are not treated comprehensively; in particular, studies with frozen midge larvae only for use to feed experimental animals are totally disregarded.

Supplements to Current Bibliography 2007

- Anjos, A. F. 2007a. *Influência das fases hídricas e tipos de substratos artificiais sobre larvas de Chironomidae no trecho superior do rio Paraná*. - Diss., Univ. Estad. Maringá.
- Anonymus 2007a. Ernst Josef Fittkau turns 80. - *Ecotropica* 13: 80.
- Arimoro, F. O., Ikomi, R. B. and Efemuna, E. 2007a. Macroinvertebrate community patterns and diversity in relation to water quality status of River Ase, Niger Delta, Nigeria. - *J. Fish. aquat. Sci.* 2: 337-344.
- Axford, Y. 2007a. *Interglacial temperature variability in the high latitude North Atlantic region inferred from subfossil midges, Baffin Island (Arctic Canada) and Iceland*. - Ph. D. Diss., Univ. Colo., Boulder.
- Chakrabarty, D. 2007a. Comparative utilization of phosphorus from sedimentary and igneous phosphate rock by major biotic components of aquatic ecosystem. - *Int. J. envir. Sci. Technol.* 4: 43-48.
- Corrêa, M. O. D. A. 2007a. *Estudo da fauna aquática associada à vegetação do mangue do Rio da Fazenda, Ubatuba (SP) através de experimentos de manipulação*. - Diss., Univ. Estad. Paulista, Botucatu. v + 122 pp.
- Coulson, S. J. 2007a. Terrestrial and freshwater invertebrate fauna of the High Arctic archipelago of Svalbard. - *Zootaxa* 1448: 41-68.
- Fang, J.-F., Lei, X.-L., Cao, T.-H. and Zong, Z.-Sh. 2007a. (Influence of *Chironomus dorsalis* Meigen larva life by environment factor and study of killing the larva) - *JingShui Jishu* (= *Water Purific. Technol.*) 26:1-3.
- Filippi, M.-A., Heiri, O., Arpent, E., Angeli, N., Bortolotti, M., Lotter, A. F. e van der Borg, K. (2005) 2007a. Studio paleolimnológico del Lago Nero di Cornisello (Parco Naturale Adamello-Brenta, Trentino). - *Studi trent. Sci. nat., Acta geol.* 82: 261-278.
- Filippi, M.-A., Heiri, O., Arpent, E., Angeli, N., Bortolotti, M., Lotter, A. F. e van der Borg, K. (2005) 2007b. Evoluzione paleoambientale dal Tardoglaciale a oggi ricostruita attraverso lo studio dei sedimenti del Lago di Lavarone (Altopiano di Folgaria e Lavarone, Trentino). - *Studi trent. Sci. nat., Acta geol.* 82: 279-298.
- Gilka, W., Kłonowska-Olejnik, M. and Godunko, R. J. 2007a. On the biology of *Symbiocladius rhithrogenae* (Zavřel, 1924) (Diptera: Chironomidae) from the Chornohora Mts., Ukraine. - *Pol. J. Ent.* 76: 285-291.
- Gritzalis, K. C., Karaouzas, I. and Skoulikidis, N. 2007a. Assessing the ecological quality of running waters of Thrace region (NE Greece) by the use of macroinvertebrate indicators. - *Fresenius envir. Bull.* 15: 1182-1188.
- Guo, Y. H. and Wang, X. H. 2007a. *Zavrelia bragremia* [for: *bragremia*] sp. nov. from China (Diptera, Hironomidae [sic!], Tanytarsini). - *Dong Wu Fen Lei Xue Bao* (= *Acta zootaxon. sin.*) 32: 318-320.
- Guo, Y. H. and Wang, X. H. 2007b. *Stempelinella depilisa* sp. nov. from China (Diptera, Chironomidae, Tanytarsini). - *Dong*

- Wu Fen Lei Xue Bao (= *Acta zootaxon. sin.*) 32: 321-323.
- Guo, Y. H. and Wang, X. H. 2007c. (A new combination *Cladotanytarsus gracilistylus* (Chaudhuri & Datta, 1992) (Diptera: Chironomidae) from China.) - *Kun Chong Fen Lei Xue Bao* (= *Entomotaxonomia*) 29: 297-300.
- Heiri, O., Filippi, M.-L. and Lotter, A. F. (2005) 2007a. Lateglacial summer temperature in the Trentino area (Northern Italy) as reconstructed by fossil chironomid assemblages in Lago di Lavarone (1100 m a.s.l.) - *Studi trent. Sci. nat., Acta geol.* 82: 299-308.
- Int Panis, L. 2007a. Genetische variatie in populaties van *Chironomus nuditarsis* Keyl (Diptera, Chironomidae); implicaties voor biodiversiteit. (Genetic variation in populations of *Chironomus nuditarsis* Keyl (Diptera, Chironomidae); implications for biodiversity.) - In: *Congr. Watersysteemkennis 2006-2007, Aquatische Biodiversiteit. Water 2007/IMS/P/0089*: 3 pp.
- Janke, H. and Trivinho-Strixino, S. 2007a. Colonization of leaf litter by aquatic macroinvertebrates: a study in a low order tropical stream. - *Acta limnol. bras.* 19: 109-115.
- Karaouzas, I., Gritzalis, K. C. and Skoulikidis, N. 2007a. Land use effects on macroinvertebrate assemblages and stream quality along an agricultural river basin. - *Fresenius envir. Bull.* 16: 645-653.
- Kerckhof, F., Haelters, J. and Gollasch, S. 2007a. Alien species in the marine and brackish ecosystem: the situation in Belgian waters. - *Aquat. Invasions* 2: 243-257.
- Lei, P., Liu, L.-j., Zhang, J.-s. and Chen, Y.-q. 2007a. (A study on microbiological control on chironomid pollution in city water supply) - *Wei Sheng Wu Xue Tong Bao* (= *Microbiology*) 34: 296-299.
- Lin, H.-b., Song, Zh.-hui and Wang, F.-f. 2007a. (Toxic effect of triphenyltin on *Chironomus plumosus* larve, *Galba perversa* and *Tubifex tubifex*) - *Qingdao Keji Daxue Xuebao* (= *J. Qingdao Univ. Sci. Technol., nat. Sci. Edn*) 28: 386-389, 393.
- Liu, L.-j., Zhang, J.-s., Chen, Y.-q. and Zhou, L. 2007a. (Growth, reproduction regularity and prediction of chironomid midge in water supply system) - *Gong Shui Ji Shu* (= *Water Technol.*) 1, 2:10-13.
- Michailova, P., Sella, G. and Petrova, N. 2007a. Effect of heavy metals on the genome of a model insect group (Diptera, Chironomidae). - In: *Evoluytsiya i ekologiya 2007, Proc.*, pp. 36-44. Sofia.
- Mo, W.-j. 2007a. (Discussion on inactivation methods of *Chironomus* larva in water treatment process) - *Gongye Anquan Yu Huanbao* (= *Ind. Saf. envir. Protect.*) 33(5): 21-22.
- Moubayed-Breil, J. et Dia, A. (2006) 2007a. Espèces de Chironomes nouvelles pour la faune du Liban et du Proche-Orient [Diptera, Chironomidae]. - *Ephemera* 8: 101-107.
- Moubayed-Breil, J., Lounaci, A. and Lounaci-Daoudi, D. (2006) 2007a. Non-biting midges from Algeria, North Africa [Diptera, Chironomidae]. - *Ephemera* 8: 93-99.
- Murray, D. A. and Baars, J. R. 2007a. *Parapsectra uliginosa* Reiss, 1969 (Diptera, Chironomidae) new to Ireland and association of exuviae of *Procladius* Pe1 (sensu Langton) with *P. simplicistilus* Freeman, 1948. - *Dipterists Dig. Ser.* 13: 166-168.
- Niitsuma, H. 2007b. *Rheopelopia* and two new genera of Tanypodinae (Diptera, Chironomidae) from Japan. - *Jap. J. syst. Ent.* 13: 99-116.
- Pan, J., Zhang, J.-s. and Sun, T.-h. 2007a. (Control of Chironomidae larvae in water treatment by *Bacillus thuringiensis* var. *israelensis* de Bariac) - *Sheyang Shifan Daxue Xuebao-Ziranhexue Ban* (= *J. Shenyang Norm. Univ. (Nat. Sci.)*) 25: 80-82.
- Perkovsky, E. E., Rasnitsyn, A. P., Vlaskin, A. P. and Taraschuk, M. V. 2007a. A comparative analysis of the Baltic and Rovno amber arthropod faunas: representative samples. - In: Brothers, D. J. and Mostovski, M. B. (eds): *Congr. Proc. Fossils X 3, Pretoria, South Africa, 7-11 February 2005. Natal Mus., Pietermaritzburg. Afr. Invertebrates* 48: 229-245.
- Punti Casadellà, T. 2007a. *Ecologia de les comunitats de quironòmids en rius mediterranis de referència. (Ecology of Chironomidae communities in Mediterranean reference streams.)* - Tes. doct., Univ. Barcelona. xx + 151 pp.
- Resende, D. L. M. C. e Takeda, A. M. 2007a. Larvas de Chironomidae (Diptera) em três reservatórios do Estado do Paraná, Brasil. - *Revta bras. Zool.* 9: 167-176.

- Rieradevall, M., Chaves, M. L. and Prat, N. 2007a. High altitude Chironomidae (Diptera) of Serra da Estrela (Portugal): additions to the Portuguese and Iberian Peninsula fauna. - *Graellsia* 63: 273-278.
- Rodríguez-Barrios, J., Ospina-Torres, R., Gutiérrez, J. D. y Ovalle, H. 2007a. Densidad y biomasa de macroinvertebrados acuáticos derivantes en una quebrada tropical de montaña (Bogotá, Colombia). - *Caldasia* 29: 397-412.
- Rosin, G. C. 2007a. *Larvas de Chironomidae em substratos artificiais: comparações e padrões de colonização em três rios de planície de inundação do alto rio Paraná, Brasil*. - Mast. Thes., Univ. Estad. Maringá.
- Shitikov, V. K., Zincheko, T. D. i Golovatyuk, L. V. 2007a. Struktura bazy dannykh i algoritmy obrabotki informatsii. (Structure of data base and algorithm of information processing.) - In: Bukharin, O. V. i Rozenberg, G. S. (eds.): *Bioindikatsiya ekologicheskogo sostoyania ravninnykh rek. (Bioindication of ecological state of lowland rivers)*, pp. 86-103. Nauka, Moskva.
- Sun, X.-B., Cui, F.-Y., Zhang, J.-S., Xu, F. and Liu, L.-J. 2007a. Inactivation of chironomid larvae with chlorine dioxide. - *J. hazard. Mater.* 142: 348-353.
- Taşdemir, A., Ustaoglu, M. R. ve Balık, S. 2007a. Akgöl ve Gebekirse Gölü'nün (Selçuk-İzmir) Chironomidae (Diptera-Arthropoda) Faunası. (Chironomidae (Diptera - Arthropoda) fauna of Akgöl and Lake Gebekirse (Selçuk-İzmir). - *Ulusal Su Günleri 2007 Sempozyumu, 16-18 Mayıs 2007, Antalya, Türk Sucul Yaşam Dergisi*, pp. 224-228.
- Veltz, I., Azar, D. and Nel, A. 2007a. New chironomid flies in Early Cretaceous Lebanese amber (Diptera: Chironomidae). - In: Brothers, D. J. and Mostovski, M. B. (eds.): *Congr. Proc. Fossils X 3, Pretoria, South Africa, 7-11 February 2005. Natal Mus., Pietermaritzburg*, pp. 169-191. [also published as: *Afr. Invert.* 48: 169-191.]
- Vidinova, Y. N., Yaneva, I. Y., Tyufekchieva, V. G. and Zadneprovski, B. E. 2007a. Composition and abundance of macroinvertebrate communities in the Arda River (South Bulgaria) in relation to some ecological factors. - *Acta zool. bulg.* 59: 309-324.
- Walker, I. 2007a. The macrofauna associated with *Eichhornia crassipes* roots during the phase of declining eutrophication in the Balbina Reservoir (Presidente Figueiredo, Amazonas, Brasil). - *Acta limnol. bras.* 19: 473-483.
- Werner, D., Reusch, W. and Pont, A. C. 2007a. *Dicranomyia didyma* (Meigen) (Diptera: Limoniidae) as a larval predator of Chironomidae and Simuliidae and as adult prey of Muscidae (Diptera), with notes on the developmental cycle. - *Entomologist's mon. Mag.* 143: 101-111.
- Würdig, N. L., Cenzano, C. S. S. and Motta Marques, D. 2007a. Macroinvertebrate communities structure in different environments of the Taim Hydrological System in the state of Rio Grande do Sul, Brazil. - *Acta limnol. bras.* 19: 427-438.
- Zhang, Ch.-h., Zhu, Sh.-q., Chen, Y., Qi, Ch.-d., Ji, Y., Zhu, Y.-j. and Zhao, J. 2007a. (Analysis of water quality in Miyun reservoir) - *Huanjing Kexue Yu Jishu (= Envir. Sci. Technol.)* 2007(z1): 63-64, 96.
- Zhang, R.-L. and Wang, X.-H. 2007a. A new species of subgenus *Polypedilum* (s. str.) Kieffer (Diptera: Chironomidae) from China. - *Entomotaxonomia* 29: 293-296.
- Zhang, R.-L. and Wang, X.-H. 2007b. Two new species of *Polypedilum* (*Tripodura*) Townes from oriental China (Diptera, Chironomidae). - *Dong Wu Fen Lei Xue Bao (= Acta zootaxon. sin.)* 32: 787-791.
- Zhu, D., Su, H.-y. and Zheng, Y.-ch. 2007a. (A review on infesting and bio-prevention of chironomid larva pollution in municipal water system) - *Chuxiong Shifan Xueyuan Xuebao (= J. Chuxiong Normal Univ.)* 22, 6: 43-47.
- Zhu, G.-q. 2007a (A study on prevention of chironomid larvae pollution of secondary water supply) - *Guangdong Keji (= Guangdong Sc. Technol.)* 164: 36-37.
- Zinchenko, T. D. i Moller Pilot, Kh. K. M. 2007a. Indikatsionnaya rol' ekzuviev kukolok khironomid pri otsenke kachestva vodnykh ekosistem ravninnykh rek. (Indicator role of chironomid pupal exuviae in evaluation of the quality of aquatic ecosystems of lowland rivers.) - In: Bukharin, O. V. i Rozenberg, G. S. (eds.): *Bioindikatsiya ekologicheskogo sostoyania ravninnykh rek. (Bioindication of ecological state of lowland rivers)*, pp. 237-242. Nauka, Moskva.
- Zinchenko, T. D. i Rozenberg, G. S. 2007a. Otsenka kachestva bioindikatorov.

- (Evaluation of the quality of bioindicators.) - In: Bukharin, O. V. i Rozenberg, G. S. (eds.): *Bioindikatsiya ekologicheskogo sostoyania ravninnykh rek. (Bioindication of ecological state of lowland rivers)*, pp. 370-380. Nauka, Moskva.
- Zinchenko, T. D., Golovatyuk, L. V. i Zagorskaya, E. P. 2007a. Strukturnaya organizatsiya soobshchestv makrozoobentosa ravninnykh rek pri antropogennom vozdeistvii. (Structural organization of macrozoobenthos communities of lowland rivers under anthropogenic influence. - In: Bukharin, O. V. i Rozenberg, G. S. (eds.): *Bioindikatsiya ekologicheskogo sostoyania ravninnykh rek. (Bioindication of ecological state of lowland rivers)*, pp. 217-233. Nauka, Moskva.
- Zinchenko, T. D., Sharyi, P. A., Sharaya, L. S., Shitikov, V. K. i Abrosimova, E. V. 2007a. Znachenie rel'efa v otsenke gidrobiologicheskogo sostoyaniya malykh rek. (Importance of the relief in evaluation of the hydrobiological state of small rivers.) - In: Bukharin, O. V. i Rozenberg, G. S. (eds.): *Bioindikatsiya ekologicheskogo sostoyania ravninnykh rek.*
- Supplements to Current Bibliography 2008**
- Al-Saffar, M. A. T. 2008a. An invitation for stopping cholera in Iraq before its outbreak. - *Nature Iraq Paper 2008*: 5 pp. Sulaimani, Kurdistan, Iraq.
- Amirkolaie, A. K. 2008a. Diet composition of sturgeon fry (*Acipenser persicus*) reared in fertilized earthen ponds. - *Aquacult. Res.* 40: 112-117.
- Andersen, T., Mendes. H. F. and de Pinho, L. C. 2008a. Two new species of *Stenochironomus* KIEFFER, 1919 from the Brazilian Atlantic Rainforest (Diptera: Chironomidae). - *Studia dipterol.* 14: 263-269.
- Armengol, X., Anton-Pardo, M., Atienzar, F., Echevarrias, J. L. and Barba, E. 2008a. Limnological variables relevant to the presence of the endangered white-headed duck in southeastern Spanish wetlands during a dry period. - *Acta zool. Acad. Scient. hung.* 54, Suppl. 1: 45-60.
- Béchar, K. M., Gillis, P. L. and Wood, C. M. 2008a. Trophic transfer of Cd from larval chironomids (*Chironomus riparius*) exposed via sediment or waterborne routes, to zebrafish (*Danio rerio*): Tissue-specific and subcellular comparisons. - *Aquat. Toxic.* 90: 310-321.
- Beketov, M. A., Schäfer, R. B., Marwitz, A., Paschke, A. and Liess, M. 2008a. Long-term stream invertebrate community alterations induced by the insecticide thiacloprid: Effect concentrations and recovery dynamics. - *Sci. tot. Envir.* 405: 96-108.
- Benbow, M. E. 2008a. Role of larval sexual dimorphism, biased sex ratios, and habitat on the energetics of a tropical chironomid. - *Envir. Ent.* 37: 1162-1173.
- Borderelle, A.-L., Gilmette, C., Lovy, C., Gerdeaux, D. and Verneaux, V. 2008a. Macroinvertebrate $\delta^{13}\text{C}$ variability analysis for the assessment of lake trophic functioning. - *Fundam. appl. Limnol.* 172: 289-300.
- Berezina, N. A. 2008a. Spatial distribution of macrofauna in a littoral zone with drifting macroalgae in the Neva estuary. - *Eston. J. Ecol.* 57: 198-213.
- Botelho, S. C., Tyagi, A., Hessle, V., Östlund Farrants, A. K. and Visa, N. 2008a. The association of Brahma with the Balbiani ring 1 gene of *Chironomus tentans* studied by immunoelectron microscopy and chromatin immunoprecipitation. - *Insect molec. Biol.* 17: 505-513.
- Brodin, Y., Lundström, J. O. and Paasivirta, L. 2008a. *Tavastia yggdrasilia*, a new orthoclad midge (Diptera: Chironomidae) from Europe. - *Aquat. Insects* 30: 261-267.
- Broza, M., Gancz, H. and Kashi, Y. 2008a. The association between non-biting midges and *Vibrio cholerae*. - *Envir. Microbiol.* 10: 3193-3200.
- Chen, Ch.-x. and Lu, P. 2008a. (Prevention of chironomid and its larvae in waterworks using advanced treatment) - *Zhongguo Jishui Panshui (= China Water and Wastewater)* 24, 6: 95-98.
- Chen, J. H., Chen, F. H., Zhang, E. L., and Zhou, A. F. 2008a. (A preliminary study on chironomid-based salinity reconstruction for Sugan Lake in the last millennium). - *Quatern. Sci.* 28: 338-344.
- Cheng, M. and Wang, X. 2008a. New species of *Clinotanypus* Kieffer, 1913 (Chironomidae: Tanypodinae) from China. - *Zootaxa* 1944: 53-65.
- Chételat, J., Amyot, M., Cloutier, L. and Poulain, A. 2008a. Metamorphosis in chironomids,

- more than mercury supply, controls methylmercury transfer to fish in high Arctic lakes. - *Envir. Sci. Technol.* 42: 9110-9115.
- Clayton, J. A. and Westbrook, C. J. 2008a. The effect of the Grand Ditch on the abundance of benthic invertebrates in the Colorado River, Rocky Mountain National Park. - *River Res. Applics* 24: 975-987.
- Copp, G. H., Kováč, V., Zweimüller, I., Dias, A., Nascimento, M. and Balázová; M. 2008a. Preliminary study of dietary interactions between invading Ponto-Caspian gobies and some native fish species in the River Danube near Bratislava (Slovakia). - *Aquat. Invasions* 3: 193-200.
- Corbi, J. J. and Trivinho-Strixino, S. 2008a. Relationship between sugar cane cultivation and stream macroinvertebrate communities. - *Braz. Archs Biol. Tech.* 51: 569-579.
- Cremona, F., Planas, D. and Lucotte, M. 2008a. Assessing the importance of macroinvertebrate trophic dead ends in the lower transfer of methylmercury in littoral food webs. - *Can. J. Fish. aquat. Sci.* 65: 2043-2052.
- Cui, Y. and Wang, H. 2008a. (Ecology of macrozoobenthic communities in two plateau lakes of Southwest China.) - *Chin. J. Oceanol. Limnol.* 26: 345-352.
- Deak, Cs., Gor, D. and Lakatos, Gy. 2008a. Phytophilous macrofauna assessments in an important aquatic bird area: the Kis-Balaton protection system. - *Acta zool. Acad. Scient. hung.* 54, Suppl. 1: 75-83.
- De Haas, E. M. and Kraak, M. H. S. 2008a. Species-specific responses of two benthic invertebrates explain their distribution along environmental gradients in freshwater habitats. - *Sci. tot. Envir.* 406: 430-435.
- De Jonge, M., van de Vijver, B., Blust, R. and Bervoets, L. 2008a. Responses of aquatic organisms to metal pollution in a lowland river in Flanders: A comparison of diatoms and macroinvertebrates. - *Sci. tot. Envir.* 407: 615-629.
- De Lange, H. J., Griethuysen, C. van and Koelmans, A. A. 2008a. Sampling method, storage and pretreatment of sediment affect AVS concentrations with consequences for bioassay responses. - *Envir. Pollut.* 151: 243-251.
- De Lange, H. J., Peeters, E. T. H. M., Harmsen, J., Maas, H. and De Jonge, J. 2008a. Seasonal variation of total and biochemically available concentrations of PAHs in a floodplain lake sediment has no effect on the benthic invertebrate community. - *Chemosphere* 75: 319-326.
- De Sousa, S., Pinel-Alloul, B. and Cattaneo, A. 2008a. Response of littoral macroinvertebrate communities on rocks and sediments to lake residential development. - *Can. J. Fish. aquat. Sci.* 65: 1206-1216.
- Donato, M., Massaferrero, J. and Brooks, S. J. 2008a. Chironomid (Chironomidae: Diptera) checklist from Nahuel Huapi National Park, Patagonia, Argentina. - *Revta Soc. ent. argent.* 67: 163-170.
- Dussault, E. B., Balakrishnan, V. K., Solomon, K. R. and Sibley, P. K. 2008a. Chronic toxicity of the synthetic hormone 17 α -ethinylestradiol to *Chironomus tentans* and *Hyalella azteca*. - *Envir. Toxic. Chem.* 27: 2521-2529.
- Eagles-Smith, C. A., Suchanek, T. H., Colwell, A. E. and Anderson, N. L. 2008a. Mercury trophic transfer in a eutrophic lake: the importance of habitat-specific foraging. - *Ecol. Applics* 18, spec. Issue: A196-A212.
- Eggermont, H., Kennedy, D., Hasiotis, S. T., Verschuren, D. and Cohen, A. 2008a. Distribution of living larval Chironomidae (Insecta: Diptera) along a depth transect at Kigoma Bay, Lake Tanganyika: implications for palaeoenvironmental reconstruction. - *Afr. Ent.* 16: 162-184.
- Eggermont, H., Verschuren, D., Fagot, M., Rumes, B., Van Bocxlaer, B. and Kröpelin, S. 2008a. Aquatic community response in a groundwater-fed desert lake to Holocene desiccation of the Sahara. - *Quatern. Sci. Rev.* 27: 2411-2425.
- Fernández, H. R., Romero, F. and Domínguez, E. 2009a. Intermountain basins use in subtropical regions and their influences on benthic fauna. - *River Res. Applics* 25: 181-193.
- Filippi, M. L., Angeli, N., Marziali, L., Matouk, A. H. I., Bertuzzi, E., Lencioni, V. and Cantonati, M. 2008a. Paleolimnological techniques applied to spring habitats: Challenges and limits as highlighted by the CRENO DAT project (southeastern Alps, Italy). - *Verh. int. Verein. Limnol.* 30: 329-332.
- Fleissner, Ge., Fleissner, Gu., Kaiser, T., Schuchardt, K. and Neumann, D. 2008a. Multiple extraretinal photoreceptors in the

- intertidal midge *Clunio marinus*. - In: Warrant, E. and Nilsson, D. E. (eds.): *2nd Int. Conf. Invert. Vision*, p. 174. Lund Univ.
- Fleissner, Gu., Fleissner, Ge., Schuchardt, K. and Neumann, D. 2008a. Larval ocelli as a putative moonlight receptor. - In: Warrant, E. and Nilsson, D. E. (eds.): *2nd Int. Conf. Invert. Vision*, p. 62. Lund Univ.
- Fu, Y. and Wang, X.-H. 2008a. A new species of *Paratrichocladius* Santos Abreu (Diptera: Chironomidae) from China. - *Si Chuan Dong Wu* (= *Sichuan J. Zool.*) 27: 728-730.
- Fusari, L. M., Roque, F. O. and Hamada, N. 2008a. Sponge-dwelling chironomids in the Upper Paraná River (Brazil): Little known but potentially threatened species. - *Neotrop. Ent.* 37: 522-527.
- Gardarsson, A., Petersen, A. and Einarsson, A. 2008a. Population limitation in dabbling ducks at Myvatn, Iceland. - *Acta zool. Acad. Scient. hung.* 54, Suppl. 1: 131-143.
- Gavrilă, L., Burlibaşa, L., Uşurelu, M. D., Radu, I., Magdalena, L. M., Ardelean, A. and Cărăbaş, M. 2008a. Chromosomal rearrangements in *Chironomus* sp. as genosensors for monitoring environmental pollution. - *Rom. biotechnol. Lett.* 13: 3962-3969.
- Gergs, R. and Rothhaupt, K.-O. 2008a. Feeding rates, assimilation efficiencies and growth of two amphipod species on biodeposited material from zebra mussels. - *Freshwat. Biol.* 53: 2494-2503.
- Gilka, W. 2008a. An intraspecific morphological variability of *Zavrelia pentatoma* KIEFFER (Diptera: Chironomidae). - *Dipteron* 24: 11-15.
- Golygina, V. V. and Kiknadze, I. I. 2008a. The revision of chromosome I (AB) mapping in *Chironomus plumosus* group (Diptera: Chironomidae). - *Comp. Cytogenet.* 2: 37-55.
- Goudie, R. I. and Gilliland, S. G. 2008a. Aspects of distribution and ecology of harlequin ducks on the Torrent River, Newfoundland. - *Waterbirds* 31, spec. Issue 2: 92-103.
- Grimaldi, D. A. and Triplehorn, D. M. 2008a. Insects from the Upper Miocene Grubstake Formation of Alaska. - *Am. Mus. Novit.* 3612: 1-19.
- Gunderina, L. I., Kiknadze, I. I., Mikhailova, P. V. i Krastanov, B. D. 2008a. Izmenchivost' i divergenziya genomnykh DNK *Chironomus plumosus* L. i *Chironomus balatonicus* Dévai, Wülker et Scholl (Diptera, Chironomidae) v prirodnykh populyatsiyakh Rossii i Bolgarii. (Variation and divergence of genomic DNA in *Chironomus plumosus* L. and *Chironomus balatonicus* Dévai, Wülker et Scholl (Diptera, Chironomidae) of natural populations in Russia and Bulgaria.) - *Evrasiat. ent. Zh.* (= *Euroas. ent. J.*) 7: 167-174.
- Gyllström, M., Lakowitz, T., Brönmark, C. and Hansson, L.-A. 2008a. Bioturbation as driver of zooplankton recruitment, biodiversity and community composition in aquatic ecosystems. - *Ecosystems* 11: 1120-1132.
- Hirabayashi, K., Tanizaki, S. and Yamamoto M. 2008a. Chironomid (Diptera, Chironomidae) fauna in a filtration plant in Japan. - In: Robinson, W. H. and Bajomi, D. (eds.): *Proc. 6th Int. Conf. Urban Pests*: 187-192. OOK-Pr. Kft., Veszprém.
- Hoffmann, A. A., Reynolds, K. T., Nash, M. A. and Weeks, A. R. 2008a. A high incidence of parthenogenesis in agricultural pests. - *Proc. R. Soc. B* 275: 2473-2481.
- Holmes, N. 2008a. Validation of chironomid-inferred temperature reconstructions in Iceland: the potential for reconstructing quantitative changes in Holocene climate. - *Geogr. helv.* 63: 4-14.
- Hughes, S. J., Ferreira, T. and Cortes, R. V. 2008a. Hierarchical spatial patterns and drivers of change in benthic macroinvertebrate communities in an intermittent Mediterranean river. - *Aquat. Conserv. mar. Freshwat. Ecosyst.* 18: 742-760.
- Ilkova, J., Michailova, P., Petrova, N., Sella, G., Hankeln, T. and Schmidt, E. 2008a. Repetitive DNA localization in two homosequential species of the genus *Chironomus* Kieffer (Diptera: Chironomidae) and their genome reaction to anthropogenic factors. - *Acta zool. bulg., Suppl.* 2: 41-48.
- Inoue, E., Kimura, G. and Hirabayashi, K. 2008a. Chironomids (Diptera: Chironomidae) attracted to vending machines in the middle reach of the Shinano River, Japan. - In: Robinson, W. H. and Bajomi, D. (eds.): *Proc. 6th Int. Conf. Urban Pests*: 177-185. OOK-Pr. Kft., Veszprém.
- Jacobsen, D. and Marín, R. 2008a. Bolivian Altiplano streams with low richness of macroinvertebrates and large diel fluctuations in temperature and dissolved oxygen. - *Aquat. Ecol.* 42: 643-656.

- Jacobsen, R. E. 2008a. A key to the pupal exuviae of the midges (Diptera: Chironomidae) of Everglades National Park, Florida. - *U. S. Geol. Surv. Scient. Invest. Rep. 2008-5082*: 119 pp.
- Janas, U. and Barańska, A. 2008a. What is the diet of *Palaemon elegans* Rathke, 1837 (Crustacea, Decapoda), a non-indigenous species in the Gulf of Gdańsk (southern Baltic Sea)? - *Oceanologia* 50: 221-237.
- Jarzebowski, E. A., Azar, D. and Nel, A. 2008a. A new chironomid (Insecta: Diptera) from Wealden amber (Lower Cretaceous) of the Isle of Wight (UK). - *Geol. Acta* 6: 285-291.
- Javanshir, A., Shapoori, M. and Jamili, S. 2008a. Diversity of benthic invertebrate fauna and secondary production in southern Caspian Sea basin, case study on Tajan River estuary. - *J. Fish. aquat. Sci.* 3: 353-365.
- Jiang, J., Wen, F.-n., Su, H.-w., Li, X.-y., He, P. and Yan, Y.-j. 2008a. (Population dynamics, annual production, and trophic basis of dominant chironomids in Tanqigou Stream.) - *Sheng Tai Xue Za Zhi (= Chin. J. Ecol.)* 27: 1503-1509.
- Johnson, A. A. and Kleve, M. G. 2008a. A new species of mermithid (Nematoda) in chironomids eclosing from a northern Minnesota bog. - *J. Parasit.* 94: 1342-1345.
- Johnson, J. H., McKenna Jr., J. E., Nack, C. C. and Chalupnicki, M. A. 2008a. Diel diet composition and feeding activity of round goby in the nearshore region of Lake Ontario. - *J. Freshwat. Ecol.* 23: 607-612.
- Jorcín, A. and Nogueira, M. G. 2008a. Benthic macroinvertebrates in the Paranapanema reservoir cascade (southeast Brazil). - *Braz. J. Biol.* 68, 4, Suppl.: 1013-1024.
- Jung, S. W., Nguyen, V. V., Nguyen, Q. H. and Bae, Y. J. 2008a. Aquatic insect faunas and communities of a mountain stream in Sapa Highland, northern Vietnam. - *Limnology* 9: 219-229.
- Kibichii, S., Shivoga, W. A., Muchiri, M., Enanga, E. and Miller, S. N. 2008a. Seasonality in water quality and its influence on the abundance and distribution of phytoplankton and chironomid larvae in Lake Nakuru, Kenya. - *Verh. int. Verein. Limnol.* 30, 3: 333-338.
- Kiknadze, I. I., Golygina, V. V., Broshkov, A. D., Gunderina, L. I. and Istomina, A. G. 2008a. Mystery of *Chironomus dorsalis* Meigen karyotype (Diptera: Chironomidae). - *Comp. Cytogenet.* 2: 21-35.
- Koščo, J., Manko, P., Miklisová, D., Košuthová, L. 2008a. Feeding ecology of invasive *Percottus glenii* (Perciformes, Odontobutidae) in Slovakia. - *Czech J. Anim. Sci.* 53: 479-486.
- Krakowiak, P. J. and Pennuto, C. M. 2008a. Fish and macroinvertebrate communities in tributary streams of eastern Lake Erie with and without round gobies (*Neogobius melanostomus*, Pallas 1814). - *J. Gt. Lakes Res.* 34: 675-689.
- Krashennikov, A. B. i Makarchenko, E. A. 2008a. *Prosmittia verae* sp. n. - novyi vid komarov-zvontsov (Diptera: Chironomidae: Orthoclaadiinae) iz okrestnostei Sochi (Severnyi Kavkaz). (*Prosmittia verae* sp. n. - a new chironomid species (Diptera: Chironomidae: Orthoclaadiinae) from the environs of Sochi Town (the Northern Caucasus)). - *Kavkaz. ent. Byul.* 4: 359-361.
- Krastanov, B. D. and Michailova, P. V. 2008a. Cytotaxonomic characteristic of four species of *plumosus* group in genus *Chironomus* Meigen 1803 (Diptera: Chironomidae) from Bulgaria and Poland. - *Acta zool. bulg., Suppl.* 2: 49-60.
- Krivosheina, M. G. 2008a. O pitanii nasekomykh tsianobakteriyami. (On insect feeding on cyanobacteria.) - *Paleont. Zh.* 2008, 6: 26-29. [also published in: *Paleont. J.* 42: 596-599.]
- Landeiro, V. L., Hamada, N. and Melo, A. S. 2008a. Responses of aquatic invertebrate assemblages and leaf breakdown to macroconsumer exclusion in Amazonian "terra firme" streams. - *Fundam. appl. Limnol.* 172: 49-58.
- Larsen, S., Vaughan, I. P. and Ormerod, S. J. 2008a. Scale-dependent effects of fine sediments on temperate headwater invertebrates. - *Freshwat. Biol.* 54: 203-219.
- Lerner, A., Meltser, N., Sapir, N., Erlick, C., Shashar, N. and Broza, M. 2008a. Reflected polarization guides chironomid females to oviposition sites. - *J. exp. Biol.* 211: 3536-3543.
- Li, A. O. Y. and Dudgeon, D. 2008a. Food resources of shredders and other benthic macroinvertebrates in relation to shading conditions in tropical Hong Kong streams. - *Freshwat. Biol.* 53: 2011-2025.

- Lods-Crozet, B., Reymond, O. et Strawczynski, A. 2008a. Evaluation de l'état écologique de deux lacs sub-alpins suisses (canton de Vaud). - *Bull. Soc. vaud. Sci. nat.* 91: 157-173.
- Love, J. W., Taylor, C. M. and Warren Jr., M. P. 2008a. Effects of summer drought on fish and macroinvertebrate assemblage properties in upland Ouachita Mountain streams, USA. - *Am. Midl. Nat.* 160: 265-277.
- Makarchenko, E. A. and Makarchenko, M. A. 2008a. *Doithrix doriceni* sp. n. (Diptera, Chironomidae, Orthoclaadiinae) from South Primorye of the Russian Far East. - *Evraz. ent. Zh.* (= *Euroas. ent. J.*) 7: 68-70.
- Makarchenko, E. A. and Makarchenko, M. A. 2008b. The first record of *Platysmittia* Sæther (Diptera, Chironomidae, Orthoclaadiinae) in Palaearctic region. - *Evraz. ent. Zh.* (= *Euroas. ent. J.*) 7: 177-179.
- Makarchenko, E. A. and Makarchenko, M. A. 2008c. Dopolneniya i utochneniya k faune khironomid podsemeistva Orthoclaadiinae (Diptera, Chironomidae) rossiiskogo Dal'nego Vostoka. (Additions and corrections to the Orthoclaadiinae (Diptera, Chironomidae) fauna of the Russian Far East.) - In: *Chteniya pamyati Vladimira Yakovlevicha Levanidova (Vladimir Ya. Levanidov's Biennial Memorial Meetings)* 4: 172-186. Dal'nauka, Vladivostok.
- Makarchenko, E. A. and Makarchenko, M. A. 2008d. Novye nakhodki khironomid (Diptera, Chironomidae) na Dal'nego Vostoke i sopredel'nykh territoriyakh. V. Podsemeistvo Orthoclaadiinae. (New findings of chironomids (Diptera, Chironomidae) in Far East and bordering territories. V. Subfamily Orthoclaadiinae.) - In: *Chteniya pamyati Vladimira Yakovlevicha Levanidova (Vladimir Ya. Levanidov's Biennial Memorial Meetings)* 4: 187-195. Dal'nauka, Vladivostok.
- Makarchenko, E. A. and Makarchenko, M. A. 2008e. Review of the genus *Pseudosmittia* Edwards (Diptera: Chironomidae: Orthoclaadiinae) from the Russian Far East (Diptera: Chironomidae: Orthoclaadiinae). - *Russ. ent. J.* 17: 215-226.
- Makarchenko, E. A. and Yavorskaya, N. M. 2008a. *Monodiamesa kamora* sp. n. (Diptera, Chironomidae, Prodiamesinae) from the Amur River basin. - *Evraz. ent. Zh.* (= *Euroas. ent. J.*) 7: 175-176.
- Makarchenko, E. A., Wu, J. and Wang, X. 2008a. Two new species of chironomids (Diptera: Chironomidae) from China. - *Russ. ent. J.* 17: 321-323.
- Marques, T. S., Ferronato, B. O., Guardia, I., Longo, A. L. B., Trivinho-Strixino, S., Bertoluci, J. e Verdade, L. M. 2008a. Primeiro registro de larvas de *Chironomus inquinatus* Correia, Trivinho-Strixino & Michailova (Diptera, Chironomidae) vivendo no casco do cágado *Phrynops geoffroanus* Schweigger (Testudines, Chelidae) na região Neotropical. - *Biota neotrop.* 8: 201-203.
- Martínez-Guitarte, J. L., Díez, J. L. and Morcillo, G. 2008a. Transcription and activation under environmental stress of the complex telomeric repeats of *Chironomus thummi*. - *Chromosome Res.* 16: 1085-1096.
- Martinez-Guitarte, J. L., Planello, R. and Morcillo, G. 2008a. Changes in the ecdysone response pathway after exposure to different endocrine-disrupting chemicals in the aquatic larvae of *Chironomus riparius*. - *Comp. Biochem. Physiol., Pt A, -molec. integr. Physiol.* 151, Suppl.: S6-S7.
- Martinez-Guitarte, J. L., Planello, R. and Morcillo, G. 2008b. Changes in the ecdysone response pathway after exposure to cadmium, pentachlorophenol and bisphenol - A in the aquatic larvae of *Chironomus*. - *Toxic. Lett.* 180, Suppl. 1: S190-S191.
- Maul, J. D., Brennan, A. A., Harwood, A. D. and Lydy, M. J. 2008a. Effect of sediment-associated pyrethroids, fipronil, and metabolites on *Chironomus tentans* growth rate, body mass, condition index, immobilization, and survival. - *Envir. Toxic. Chem.* 27: 2582-2590.
- McIntosh, M. D., Schmitz, J. A., Benbow, M. E. and Burky, A. J. 2008a. Structural and functional changes of tropical riffle macroinvertebrate communities associated with stream flow withdrawal. - *River Res. Applics* 24: 1045-1055.
- McLachlan, A. J., Pike, T. W. and Thomason, J. C. 2008a. Another kind of symmetry: Are there adaptive benefits to the arrangement of mites on an insect host? - *Ethol. Ecol. Evolut.* 20: 257-270.
- Mellon, C. D., Wipfli, M. S. and Li, J. L. 2008a. Effects of forest fire on headwater stream macroinvertebrate communities in eastern Washington, U.S.A. - *Freshwat. Biol.* 53: 2331-2343.

- Mendes, H. F. and Andersen, T. 2008a. A review of *Antillocladius* Sæther and *Litocladius* Mendes, Andersen et Sæther, with the description of two new Neotropical genera (Diptera, Chironomidae, Orthoclaadiinae). - *Zootaxa* 1887: 1-75.
- Mičetić, V., Bučar, M., Ivković, M., Piria, M., Krulik, I., Mihoci, I., Delić, A. and Kučinić, M. 2008a. Feeding ecology of *Sabanejewia balcanica* and *Cobitis elongata* in Croatia. - *Folia zool.* 57: 181-190.
- Michailova, P. and Jablonska-Brana, I. 2008a. Chromosome polymorphism of *Camptochironomus tentans* (FABRICIUS 1805) (Diptera: Chironomidae) from Łukajno (Mazurian Lakeland), Poland. - *Acta zool. bulg.* 50: 155-163.
- Michailova, P., Krastanov, B. and Matena, J. 2008a. Chromosomal polymorphism of *Chironomus plumosus* (Linnaeus, 1758) (Diptera: Chironomidae) from Bulgaria (Pazardjik region) and the Czech Republic (Ceske Budejovice region). - *Comp. Cytogenet.* 2: 57-65.
- Milesi, S. V., Biasi, C., Restello, R. M. e Hepp, L. U. 2008a. Efeito de metais Cobre (Cu) e Zinco (Zn) sobre a comunidade de macroinvertebrados bentônicos em riachos do sul do Brasil. - *Acta Scient. biol. Sci.* 30: 283-289.
- Milner, A. M., Robertson, A. L., Monaghan, K. A., Veal, A. J. and Flory, E. A. 2008a. Colonization and development of an Alaskan stream community over 28 years. - *Front. Ecol. Envir.* 6: 413-419.
- Miserendino, M. L., Brand, C. and Di Prinzio, C. Y. 2008a. Assessing urban impacts on water quality, benthic communities and fish in streams of the Andes Mountains, Patagonia (Argentina). - *Wat. Air Soil Pollut.* 194: 91-110.
- Moller Pillot, H. K.M. 2008a. Identification and ecology of the genus *Smittia* Holmgren in the Netherlands (Diptera: Chironomidae). - *Tijdschr. Ent.* 151: 245-270.
- Móra, A. and Csabai, Z. 2008a. First annotated checklist of Chironomidae of Rhodos, Greece (Insecta, Diptera). - *Spixiana* 31: 223-231
- Moubayed-Breil, J. (2007) 2008a. Non-biting midges from Continental France: new records, faunal and biogeographical outline [Diptera, Chironomidae]. - *Ephemera* 9: 17-32.
- Moubayed-Breil, J. and Langton, P. H. (2007) 2008a. Description of *Parakiefferiella normandiana* sp. n. from France [Diptera, Chironomidae]. - *Ephemera* 9: 9-16.
- Muenz, T. K., Golladay, S. W., Smith, L. L. and Vellidis, G. 2008a. Diet and abundance of Southern Two-lined Salamander larvae (*Eurycea cirrigera*) in streams within an agricultural landscape, southwest Georgia. - *SEast. Nat.* 7: 691-704.
- Nazarova, L. B., Pestryakova, L. A., Ushnitskaya, L. A. and Hubberten, H.-W. 2008a. (Chironomids (Diptera : Chironomidae) in lakes of central Yakutia and their indicative potential for paleoclimatic research.) - *Sibirsk. ekol. Zh.* 15: 141-150. [also published as: *Contemp. Probl. Ecol.* 1: 335-345.
- Negassa, A. and Prabu, P. C. 2008a. Abundance, food habits, and breeding season of exotic *Tilapia zillii* and native *Oreochromis niloticus* L. fish species in Lake Zwai, Ethiopia. - *Maejo int. J. Sci. Technol.* 2: 345-359.
- Nelson, S. M. and Thullen, J. S. 2008a. Aquatic macroinvertebrates associated with *Schoenoplectus* litter in a constructed wetland in California (USA). - *Ecol. Engng* 33: 91-101.
- Néraudeau, D., Perrichot, V., Colin, J.-P., Girard, V., Gomez, B., Guillocheau, F., Masure, E., Peyrot, D., Tostain, F., Videt, B. and Vullo, R. 2008a. A new amber deposit from the Cretaceous (uppermost Albian-lowermost Cenomanian) of southwestern France. - *Cretac. Res.* 29: 925-929.
- Nessimian, J. L., Venticinque, E. M., Zuanon, J., De Marco Jr., P., Gordo, M., Fidelis, L., D'arc Batista, J. and Juen, L. 1008a. Land use, habitat integrity, and aquatic insect assemblages in Central Amazonian streams. - *Hydrobiologia* 614: 117-131. [see: Electronic supplementary material]
- Niitsuma, H. 2008a. Review of the Japanese species of *Conchapelopia* (Insecta: Diptera: Chironomidae), with keys to the known males and pupae. - *Species Diversity* 13: 73-110.
- Obolewski, K. and Strzelczak, A. 2008a. MR&CT analysis of the qualitative and quantitative structure of macrozoobenthos in selected oxbow lakes of Northern Poland. - *Oceanol. hydrobiol. Stud.* 37: 97-104.
- Ochieng, H., de Ruyter van Steveninck, E. D. and Wanda, F. M. 2008a. Mouthpart deformities in Chironomidae (Diptera) as indicators of

- heavy metal pollution in northern Lake Victoria, Uganda. - *Afr. J. aquat. Sci.* 33: 135-142.
- Olomukoro, J. O. and Okhumale, B. O. 2008a. Differential toxic response of *Chironomus travalensis* [sic!] kieffer [sic!] (Chironomidae: Diptera) to some industrial effluents. - *Toxic. envir. Chem.* 90: 1197-1202.
- Olsson, K., Nyström, P., Stenroth, P., Nilsson, E., Svensson, M. and Granéli, W. 2008a. The influence of food quality and availability on trophic position, carbon signature, and growth rate of an omnivorous crayfish. - *Can. J. Fish. aquat. Sci.* 65: 2293-2304.
- Osczo, J., Leunda, P. M., Escala, M. C. and Miranda, R. 2008a. Summer feeding relationships of the co-occurring hatchling brown trout *Salmo trutta* and Ebro minnows *Phoxinus phoxinus* in an Iberian river. - *Acta zool. sin.* 54: 675-685.
- Ouéda, A., Guenda, W., Ouattara, A., Gourène, G., Hugueny, B. and Kabré, G. B. 2008a. Seasonal diet shift of the most important fish species in a sahelo-soudanian reservoir (Burkina Faso). - *J. Fish. aquat. Sci.* 3: 240-251.
- Oyanedel, A., Valdovinos, C., Azócar, M., Moya, C., Mancilla, G., Pedreros, P. y Figueroa, R. 2008a. Padrónes de distribución espacial de los macroinvertebrados bentónicos de la cuenca del río Aysén (Patagonia chilena). - *Gayana* 72: 241-257.
- Oyewo, E. A. and Sæther, O. A. 2008a. Revision of *Polypedilum* (*Pentapedilum*) Kieffer and *Ainuyusurika* Sasa et Shirasaki (Diptera: Chironomidae). - *Zootaxa* 1953: 1-145.
- Panizzo, V. N., Jones, V. J., Birks, H. J. B., Boyle, J. F., Brooks, S. J. and Leng, M. J. 2008a. A multiproxy palaeolimnological investigation of Holocene environmental change, between c. 10 700 and 7200 years BP, at Holebudalen, southern Norway. - *Holocene* 18: 805-817.
- Park, K. and Kwak, I.-S. 2008b. Characterization of heat shock protein 40 and 90 in *Chironomus riparius* larvae: Effects of di(2-ethylhexyl) phthalate exposure on gene expressions and mouthpart deformities. - *Chemosphere* 74: 89-95.
- Perova, S. N. 2008a. Taksonomicheskii sostav makrozoobentosa malykh karstovykh ozer Tsentral'noi Rossii. (The taxonomic composition of macrozoobenthos in central Russian small karst lakes.) - *Inland Wat. Biol.* 1: 371-379.
- Petrova, N. A. and Zhirov, S. V. 2008b. Polytene chromosomes of salivary glands of chironomids (Diptera: Chironomidae) from the Wrangel Island (Russia). - *Comp. Cytogenet.* 2: 127-130.
- Posada-García, J. A., Abril-Ramírez, G. und Parra-Sánchez, L. N. 2008a. Diversidad de los macroinvertebrados acuáticos del Páramo de Frontino (Antioquia, Colombia). - *Caldasia* 30: 441-455.
- Pozdeev, I. V. and Aleksevina, M. S. 2008a. Rol' lichinok khironomid v structure donnykh soobshchestv rek basseina Verkhnei i Srednei Kamy. (The role of chironomid larvae in the structure of river benthic communities of the Upper and Middle Kama River basins.) - *Inland Wat. Biol.* 1: 158-162.
- Principe, R. E. 2008a. Taxonomic and size structures of aquatic macroinvertebrate assemblages in different habitats of tropical streams, Costa Rica. - *Zool. Stud.* 47: 525-534.
- Principe, R. E., Boccolini, M. F. and Corigliano, M. C. 2008a. Structure and spatial-temporal dynamics of Chironomidae fauna (Diptera) in upland and lowland fluvial habitats of the Chocancharava River basin (Argentina). - *Int. Rev. Hydrobiol.* 93: 342-357.
- Proviz, V. I. 2008b. Vidoobrazovanie i khromosomnaya evolyutsiya endemichnykh khironomid roda *Sergentia* Kief. (Diptera, Chironomidae) iz ozera Baikal. Divergentsiya kariotipov i khromosomnyi polimorfizm v populyatsiyakh glubokovodnykh vidov *Sergentia nebulosa* Linevitsh et al. i *Sergentia assimilis* Proviz V. et Proviz L. (Speciation and chromosomal evolution of the Baikalian endemic chironomids of the genus *Sergentia* Kief. (Diptera, Chironomidae): Karyotype divergence and chromosomal polymorphism in the populations of deep-water species *Sergentia nebulosa* Linevitsh et al. and *Sergentia assimilis* Proviz V. et Proviz L.) - *Genetika* 44: 1627-1637. [also published in: *Russ. J. Genet.* 44: 1409-1418.]
- Pugh, P. J. A. and Convey, P. 2008a. Surviving out in the cold: Antarctic endemic invertebrates and their refugia. - *J. Biogeogr.* 35: 2176-2186.
- Puntí, T. 2008a. *Ecology of Chironomidae communities in Mediterranean reference*

- streams. - Ph. D. Thes., Univ. Barcelona. 151 pp.
- Purcell, A. H., Hoffmann, A. and Resh, V. H. 2008a. Life history of a dipteran predator (Scathophagidae: *Acanthocnema*) of insect egg masses in a northern California stream. - *Freshwat. Biol.* 53: 2426-2437.
- Qi, X., Shi, S.-D. and Wang, X.-H. 2008a. One new record of genus *Stenochironomus* Kieffer in China (Diptera, Chironomidae). - *Si Chuan Dong Wu* (= *Sichuan J. Zool.*] 27: 837-838.
- Qu, X.-D., Song, M.-Y., Park, Y.-S., Oh, Y. N. and Chon, T.-S. 2008a. Species abundance patterns of benthic macroinvertebrate communities in polluted streams. - *Annls Limnol.* 44: 119-133.
- Renouf, M. A. P. and Bell, T. 2008a. Dorset Palaeoeskimo skin processing at Phillip's Garden, Port au Choix, northwestern Newfoundland. - *Arctic* 61: 35-47.
- Richardson, J. S. 2008a. Aquatic arthropods and forestry: effects of large-scale land use on aquatic systems in Nearctic temperate regions. - *Can. Ent.* 140: 495-509.
- Ríos, B. 2008a. *Macroinvertebrate communities in an altoandine stream: Microhabitat conditions, drift dynamics, and organic matter and oviposition relevance.* - Ph. D. Thes., Univ. Barcelona. 154 pp.
- Robinson, C. T., Schmid, D., Svoboda, M. and Bernasconi, S. M. 2008a. Functional measures and food webs of high elevation springs in the Swiss alps. - *Aquat. Sci.* 70: 432-445.
- Rolland, N., Larocque, I., Francus, P., Pienitz, R. and Laperriere, L. 2008b. Evidence for a warmer period during the 12th and 13th centuries AD from chironomid assemblages in Southampton Island, Nunavut, Canada. - *Quatern. Res.* 72: 27-37.
- Roque, F. O. and Trivinho-Strixino, S. 2008a. Four new species of *Endotribelos* Grodhaus, a common fallen fruit-dwelling chironomid genus in Brazilian streams (Diptera: Chironomidae: Chironominae). - *Stud. neotrop. Fauna Envir.* 43: 191-207.
- Ross, R. M., Long, E. S. and Dropkin, D. S. 2008a. - Response of macroinvertebrate communities to remediation-simulating conditions in Pennsylvania streams influenced by acid mine drainage. - *Envir. Monit. Assess.* 145: 323-338.
- Sahuquillo, M., Miracle, M. R., Rieradevall, M. and Kornijów, R. 2008a. Macroinvertebrate assemblages on reed beds, with special attention to Chironomidae (Diptera), in Mediterranean shallow lakes. - *Limnetica* 27: 239-250.
- Sakelarieva, L., Yaneva, I., Uzunov, Y., Kumanski, K., Stoichev, S., Vidinova, Y. and Tyufekchieva, V. 2008a. Taxonomic composition and dominant structure of macrozoobenthos in the Blagoevgradska Bistritsa River. - *Acta zool. bulg., Suppl.* 2: 201-214.
- Saleh-Ahmed, R., Ismail, F. M., Abo-Ghalia, A. and El-Shenawy, N. S. 2008a. Pictorial keys of chironomid species (Order: Diptera) in El-Tall El-Keber wastewater treatment plant, Egypt. - *J. Ent.* 5: 334-355.
- Salsamendi, E., Garin, I., Almenar, D., Goiti, U., Napal, M. and Aihartza, J. 2008a. Diet and prey selection in Mehelyi's horseshoe bat *Rhinolophus mehelyi* (Chiroptera, Rhinolophidae) in the south-western Iberian Peninsula. - *Acta chiropterol.* 10: 279-286.
- Sánchez-Carmona, R., Encina, L., Rodríguez-Ruíz, A. and Rodríguez-Sánchez, V. 2008a. Age, growth and diet of the Iberian loach, *Cobitis paludica* in two different environments. - *Folia zool.* 57: 420-434.
- Sanseverino, A. M. and Nessimian, J. L. 2008b. The food of larval Chironomidae (Insecta, Diptera) in submerged litter in a forest stream of the Atlantic Forest (Rio de Janeiro, Brazil). - *Acta limnol. bras.* 20: 15-20.
- Schmidl, J., Sulzer, P. and Kitching, R. L. 2008a. The insect assemblage in water filled tree-holes in a European temperate deciduous forest: community composition reflects structural, trophic and physicochemical factors. - *Hydrobiologia* 598: 285-303.
- Schofield, K. A., Pringle, C. M., Meyer, J. L. and Rosi-Marshall, E. J. 2008a. Functional redundancy of stream macroconsumers despite differences in catchment land use. - *Freshwat. Biol.* 53: 2587-2599.
- Sehnal, F. and Sutherland, T. 2008a. Silks produced by insect labial glands. - *Prion* 2: 145-153.
- Sharley, D. J., Hoffmann, A. A. and Pettigrove, V. 2008a. Effects of sediment quality on macroinvertebrates in the Sunraysia region of the Murray-Darling Rivers, Australia. - *Envir. Pollut.* 156: 689-698.

- Shcherbina, G. Kh. 2008a. Struktura biotsenoza *Dreissena polymorpha* (Pallas) i rol' mollyuska v pitanii plotvy *Rutilus rutilus* (Linnaeus) ozero Pleshcheevo. (The structure of *Dreissena polymorpha* (Pallas) biocenosis and the role of mollusk in roach *Rutilus rutilus* (Linnaeus) feeding in Pleshcheevo Lake.) - *Inland Wat. Biol.* 1: 380-387.
- Siebrasse, J. P., Veith, R., Dobay, A., Leonhardt, H., Daneholt, B. and Kubitscheck, U. 2008a. Discontinuous movement of mRNP particles in nucleoplasmic regions devoid of chromatin. - *Proc. natn. Acad. Sci. U. S. A.* 105: 20291-20296.
- Silva, F. L., Ruiz, S. S., Bochini, G. L. and Moreira, D. C. 2008a. Functional feeding habits of Chironomidae larvae (Insecta, Diptera) in a lotic system from Midwestern region of São Paulo State, Brazil. - *Pan-Am. J. aquat. Sci.* 3: 135-141.
- Silva-da-Silva, L. R. 2008a. *Estudo taxonômico da família Chironomidae (Diptera, Nematocera) em dois pontos lóticos e lênticos do Parque Turístico Ecológico de Penedo, Itatiaia – RJ.* - Tes. Mestr., Univ. Fed. do Rio de Janeiro. 120 pp.
- Sinclair, C. S. and Gresens, S. E. 2008a. Discrimination of *Cricotopus* species (Diptera: Chironomidae) by DNA barcoding. - *Bull. ent. Res.* 98: 555-563.
- Siqueira, T., Roque, F. O. and Trivinho-Strixino, S. 2008b. Phenological patterns of neotropical lotic chironomids: Is emergence constrained by environmental factors? - *Austral Ecol.* 33: 902-910.
- Skalskaya, I. A., Bakanov, A. I. and Flerov, B. A. 2008a. (Study on zooperiphyton and zoobenthos of a small river.) - *Inland Wat. Biol.* 1: 84-92.
- Sodré, V. M. 2008a. *Study of the community of invertebrates in phytotelmata of Bromeliaceae with emphasis on Chironomidae (Insecta, Diptera) in a fragment of Atlantic Forest in Magé City, RJ.* - Tes. Mestr., Univ. Fed. São Carlos. 65 pp.
- Specziár, A. 2008a. Life history patterns of *Procladius choreus*, *Tanypus punctipennis* and *Chironomus balatonicus* in Lake Balaton. - *Annl. Limnol.* 44: 181-188.
- Staiber, W. 2008a. Centrosome hyperamplification with the formation of multiple asters and programmed chromosome inactivation in aberrant spermatocytes during male meiosis in *Acricotopus*. - *Cell Tissue Res.* 334: 81-91.
- Stribling, J. B., Jessup, B. K. and Feldman, D. L. 2008a. Precision of benthic macroinvertebrate indicators of stream condition in Montana. - *J. N. Am. benthol. Soc.* 27: 68-67.
- Stribling, J. B., Pavlik, K. L., Holdsworth, S. M. and Leppo, E. W. 2008a. Data quality, performance, and uncertainty in taxonomic identification for biological assessments. - *J. N. Am. benthol. Soc.* 27: 906-919.
- Stur, E. and Ekrem, T. 2008a. Description of the Alpine *Micropsectra oberoarensis* sp. n. with taxonomic comments on the *attenuata* group (Diptera: Chironomidae). - *Ent. fenn.* 19: 142-150.
- Suchanek, T. H., Eagles-Smith, C. A. and Harner, E. J. 2008a. Is Clear Lake methylmercury distribution decoupled from bulk mercury loading? - *Ecol. Applics* 18, spec. Issue: A107-A127.
- Suchanek, T. H., Eagles-Smith, C. A., Slotton, D. G., Harner, E. J., Adam, D. P., Colwell, A. E., Anderson, N. L. and Woodward, D. L. 2008a. Mine-derived mercury: effects on lower trophic species in Clear Lake, California. - *Ecol. Applics* 18, spec. Issue: A158-A176.
- Sugimaru, K., Kawai, K. and Imabayashi, H. 2008a. A new marine chironomid species of the genus *Tanytarsus* (Diptera: Chironomidae) from Okinawa, Japan. - *Plankton Benthos Res.* 3: 240-242.
- Sun, X.-B. and Cui, F.-Y. 2008a. Inactivation of chironomid larvae with ozone. - *Wat. Sci. Technol. Wat. Supply* 8: 355-361.
- Sun, X.-B. and Cui, F.-Y. 2008b. Inactivation of chironomid larvae with chlorine dioxide and chlorine. - *J. Donghua Univ. Engl. Ed.* 25: 361-365.
- Szalinska, E., Czaplicka-Kotas, A., Vignati, D. A. L., Ferrari, B. J. D. and Dominik, J. 2008a. Bioavailability of sedimentary chromium for chironomids in the upper Dunajec River (Southern Poland). - *Verh. int. Verein. Limnol.* 30, 4: 534-536.
- Takahashi, M. A., Higuti, J., Bagatini, Y. M., Zviejkovski, I. P. and Velho, L. F. M. 2008a. Composition and biomass of larval chironomid (Insecta, Diptera) as potential indicator of trophic conditions in southern Brazil reservoirs. - *Acta limnol. bras.* 20: 5-13.

- Tall, L., Méthot, G., Armellin, A. and Pinel-Alloul, B. 2008a. Bioassessment of benthic macroinvertebrates in wetland habitats of Lake Saint-Pierre (St. Lawrence River). - *J. Gt Lakes Res.* 34: 599-614.
- Tarkowska-Kukuryk, M. and Kornijów, R. 2008a. Influence of spatial distribution of submerged macrophytes on Chironomidae assemblages in shallow lakes. - *Pol. J. Ecol.* 56: 569-579.
- Tarkowska-Kukuryk, M. and Mieczan, T. 2008a. Diet composition of epiphytic chironomids of the *Cricotopus sylvestris* group (Diptera: Chironomidae) in a shallow hypertrophic lake. - *Aquat. Insects* 30: 285-294.
- Taşdemir, A., Ustaoglu, M. R., Balık, S. ve Sarı, H. M. 2008a. Batı Karadeniz Bölgesindeki (Türkiye) Bazı Göllerin Diptera ve Ephemeroptera Faunası. (Diptera and Ephemeroptera fauna of some lakes in Western Black Sea Region (Turkey).) - *J. Fish. Sci.com* 2: 252-260.
- Thompson, R., Kamenik, C., Schmidt, R., Pla, S., Rieradevall, M. and Catalán, J. 2008a. Testing a new multigroup inference approach to reconstructing past environmental conditions. - *J. Limnol.* 67: 155-162.
- Van de Meutter, F., Cottenie, K. and De Meester, L. 2008a. Exploring differences in macroinvertebrate communities from emergent, floating-leaved and submersed vegetation in shallow ponds. - *Fundam. appl. Limnol.* 173: 47-57.
- Verberk, W. C. E. P., Siepel, H. and Esselink, H. 2008a. Life-history strategies in freshwater macroinvertebrates. - *Freshwat. Biol.* 53: 1722-1738.
- Verberk, W. C. E. P., Siepel, H. and Esselink, H. 2008b. Applying life-history strategies for freshwater macroinvertebrates to lentic waters. - *Freshwat. Biol.* 53: 1739-1753.
- Vidinova, Y. N., Botev, I. S., Tyufekchieva, V. G., Nedyalkova, T. V., Yaneva, I. Y., Zadneprovski, B. E. and Varadinova, E. D. 2008a. Results of rapid hydrobiological monitoring of watersheds from the East- and West Aegean Sea River Basin Districts in Bulgaria. - *Acta zool. bulg. Suppl.* 2: 233-242.
- Vidinova, Y. N., Tyufekchieva, V. G., Yaneva, I. Y., Zadneprovski, B. E., Stoichev, S. A. and Kumanski, K. P. 2008a. Species composition and structure of macroinvertebrate communities in Arda River. - *Acta zool. bulg.* 60: 317-330.
- Vink, J. P. M. 2008a. The origin of speciation: Trace metal kinetics over natural water/sediment interfaces and the consequences for bioaccumulation. - *Envir. Pollut.* 157: 519-527.
- Vinogradova, E. M. 2008a. Six new species of *Polypedilum* Kieffer, 1912, from the Yucatán peninsula (Insecta, Diptera, Chironomidae). - *Spixiana* 31: 277-288.
- Wachalowicz, A., Czaplicka-Kotas, A. and Szalinska, E. 2008a. Bioavailability of chromium from bottom sediments for the larvae of the species *Chironomus riparius*. - *Ochrona środow.* 30, 3: 53-58.
- Ward, S. A. and Coburn, M. M. 2008a. Stepwise increases in maximum prey size of larval Creek Chubs, *Semotilus atromaculatus*, in an urbanized Ohio stream: - *NEast. Nat.* 15: 349-362.
- Waterkeyn, A., Grillas, P., Vanschoenwinkel, B. and Brendonck, L. 2008a. Invertebrate community patterns in Mediterranean temporary wetlands along hydroperiod and salinity gradients. - *Freshwat. Biol.* 53: 1808-1822.
- Werner, S. and Rothhaupt, K.-O. 2008a. Effects of the invasive Asian clam *Corbicula fluminea* on benthic macroinvertebrate taxa in laboratory experiments. - *Fundam. appl. Limnol.* 173: 145-152.
- Wood, P. J., Gunn, J. and Rundle, S. D. 2008a. Response of benthic cave invertebrates to organic pollution events. - *Aquat. Conserv. mar. Freshwat. Ecosyst.* 18: 909-922.
- Wu, J.-Y. and Wang, X.-H. 2008a. A first recorded species *Boreochlus longicoxalsetosus* (Diptera: Chironomidae: Podonominae) from China. - *Si Chuan Dong Wu (= Sichuan J. Zool.)* 27: 839-840.
- Yamamoto, M. 2008a. Redescription of *Mesomittia patrihortae* SÆTHER, 1985, from Japan (Diptera, Chironomidae). - *Jap. J. syst. Ent.* 14: 13-16.
- Yan, C. C. and Wang, X. H. 2008a. A study on morphological intraspecific variation of *Microchironomus tener* (Kieffer) (Diptera: Chironomidae).- - In Wang, X. H. (ed.): *Contemporary aquatic entomological study in East Asia. Proc. 3rd Int. Symp. aquat. Ent. E. Asia*, pp. 153-158. Nankai Univ. Pr., Tianjin.
- Yan, C., Jin, Z. and Wang, X. 2008a. *Cladopelma* Kieffer from the Sino-Indian Region (Diptera: Chironomidae). - *Zootaxa* 1916: 44-56.

- Yan, C., Jin, Z. and Wang, X. 2008b. *Paracladopelma* Harnisch from the Sino-Indian Region (Diptera: Chironomidae). - *Zootaxa* 1934: 1-29.
- Yan, C., Sæther, O. A., Jin, Z. and Wang, X. 2008a. Three new species of the *Harnischia* complex from the Sino-Indian Region, with a review of *Demicryptochironomus* Lenz (Diptera: Chironomidae). - *Zootaxa* 1968: 1-22.
- Yıldız, S., Taşdemir, A., Balık, S. ve Ustaoglu, M. R. 2008a. Kemer Baraj Gölü'nün (Aydın) Makrobentik (Oligochaeta, Chironomidae) Faunası. (Macrobenthic fauna (Oligochaeta, Chironomidae) of Kerner Dam Lake (Aydın).) - *J. Fish. Sci.com* 2: 457-465.
- Zhang, R. L. and Wang, X. H. 2008b. Geographic co-evolutionary study on *Polypedilum* (*Cerobregma*) Sæther & Sundal (Diptera: Chironomidae.) - In Wang, X. H. (ed.): *Contemporary aquatic entomological study in East Asia. Proc. 3rd Int. Symp. aquat. Ent. E. Asia*, pp. 158-171. Nankai Univ. Pr., Tianjin.
- Zilli, F. L., Montalto, L., Paggi, A. C. and Marchese, M. R. 2008a. Biometry and life cycle of *Chironomus calligraphus* Goeldi 1905 (Diptera, Chironomidae) in laboratory conditions. - *Interciencia* 33: 767-770.
- Current Bibliography 2009**
- Acosta, C. R. 2009a. (*Altitudinal distribution of macroinvertebrate communities and hydrochemical characterization of karstic headwaters of the altitude river Cañete (Peru).*) - Ph. D. Thes., Univ. Barcelona. 153 pp.
- Agra, A. R. and Soares, A. M. V. M. 2009a. Effects of two insecticides on survival, growth and emergence of *Chironomus riparius* Meigen. - *Bull. envir. Contam. Toxic.* 82: 501-504.
- Ali, A., Leckel Jr., R. J., Jahan, N., Al-Shami, S. A. and Rawi, C. S. MD. 2009a. Laboratory and field investigations of pestiferous Chironomidae (Diptera) in some man-made wetlands in central Florida, USA. - *J. Am. Mosquito Control Ass.* 25: 94-99.
- Allgood, D. W., Miller, D. A. and Kalcounis-Rueppell, M. C. 2009a. Influence of intensive pine management on dipteran community structure in coastal North Carolina. - *Envir. Ent.* 38: 657-666.
- Almeida, E. F., Oliveira, R. B., Mugnai, R., Nessimian, J. L. and Baptista, D. F. 2009a. Effects of small dams on the benthic community of streams in an Atlantic forest area of southeastern Brazil. - *Int. Rev. Hydrobiol.* 94: 179-193.
- Al-Shami, S. A., Che Salmah, M. R., Siti Azizah, M. N. and Abu Hassan, A. 2009a. Distribution and abundance of Chironomidae (Diptera) in a rice agroecosystem in Penang, Malaysia. - In: Hughes, S. J., Khadem, M. and Carvalho, M. A. (eds.): *Proc. 16th Int. Chironomid Symp., Bolm Mus. munic. Funchal (Hist. nat.), Suppl. 13*: 151-160.
- Álvarez, M. and Pardo, I. 2009a. Dynamics in the trophic structure of the macroinvertebrate community in a Mediterranean temporary stream. - *Aquat. Sci.* 71: 202-213.
- Amundsen, P.-A. and Knudsen, R. 2009a. Winter ecology of Arctic charr (*Salvelinus alpinus*) and brown trout (*Salmo trutta*) in a subarctic lake, Norway. - *Aquat. Ecol.* 43: 765-775.
- Andrei, A. E., Smith, L. M., Haukos, D. A., Surles, J. G. and Johnson, W. P. 2009a. Foraging ecology of migrant shorebirds in saline lakes of the Southern Great Plains. - *Waterbirds* 32: 138-148.
- Andrič, M., Massaferrero, J., Eicher, U., Ammann, B., Leuenberger, M. C., Martinčič, A., Marinova, E. and Brancelj, A. 2009a. A multiproxy Late-glacial palaeoenvironmental record from Lake Bled, Slovenia. - *Hydrobiologia* 631: 121-141.
- Angradi, T. R., Bolgrien, D. W., Jicha, T. M., Pearson, M. S., Taylor, D. L. and Hill, B. H. 2009a. Multispatial-scale variation in benthic and snag-surface macroinvertebrate assemblages in mid-continent US great rivers. - *J. N. Am. benthol. Soc.* 28: 122-141.
- Anjos, A. F. e Takeda, A.M. 2009a. Análise da dieta das larvas de 4^o estágio de *Cricotopus* sp. (Diptera: Chironomidae), em diferentes substratos artificiais e fases hídricas, no trecho superior do rio Paranà. - *Acta Scient.biol. Sci.* 31: 371-377.
- Anonymus 2009a. Anna Andreevna Linevitsh. - *Evraziat. ent. Zh. (= Euroas. ent. J.)* 8, Suppl. 1: 3-5.
- Arimoro, F. O. and Ikomi, R. B. 2009a. Ecological integrity of upper Warri River, Niger Delta using aquatic insects as bioindicators. - *Ecol. Indicators* 9: 455-461.

- Axford, Y., Briner, J. P., Miller, G. H. and Francis, D. R. 2009a. Paleocological evidence for abrupt cold reversals during peak Holocene warmth on Baffin Island, Arctic Canada. - *Quatern. Res.* 71: 142-149.
- Axford, Y., Geirsdóttir, Á., Miller, G. H. and Langdon, P. G. 2009a. Climate of the Little Ice Age and the past 2000 years in northeast Iceland inferred from chironomids and other lake sediment proxies. - *J. Paleolimnol.* 41: 7-24.
- Babrekar, A. A. 2009a. *The role of iron in the homeostasis & abiotic stress response in Chironomus ramosus.* - Ph. D. Thes., Univ. Pune. 117 pp.
- Baldigo, B. P., Lawrence, G. B., Bode, R. W., Simonin, H. A., Roy, K. M. and Smith, A. J. 2009a. Impacts of acidification on macroinvertebrate communities in streams of the western Adirondack Mountains, New York, USA. - *Ecol. Indicators* 9: 226-239.
- Barquín, J. and Death, R. G. 2009a. Physical and chemical differences in karst springs of Cantabria, northern Spain: do invertebrate communities correspond? - *Aquat. Ecol.* 43: 445-455.
- Barriga, J. P. and Battini, M. A. 2009a. Ecological significances of ontogenetic shifts in the stream-dwelling catfish, *Hatcheria macraei* (Siluriformes, Trichomycteridae), in a Patagonian river. - *Ecol. Freshwat. Fish* 18: 395-405.
- Benoit, J. B., Lopez-Martinez, G., Elnitsky, M. A., Lee Jr., R. E. and Denlinger, D. L. 2009a. Dehydration-induced cross tolerance of *Belgica antarctica* larvae to cold and heat is facilitated by trehalose accumulation. - *Comp. Biochem. Physiol. - A molec. integrat. Physiol.* 152: 518-523.
- Benstead, J. P., March, J. G., Pringle, C. M., Ewel, K. C. and Short, J. W. 2009a. Biodiversity and ecosystem function in species-poor communities: community structure and leaf litter breakdown in a Pacific island stream. - *J. N. Am. benthol. Soc.* 28: 454-465.
- Bertrand, K. N., Gido, K. B., Dodds, W. K., Murdock, J. N. and Whiles, M. R. 2009a. Disturbance frequency and functional identity mediate ecosystem processes in prairie streams. - *Oikos* 118: 917-933.
- Bickel, T. O. and Closs, G. P. 2009a. Impact of partial removal of the invasive macrophyte *Lagarosiphon major* (Hydrocharitaceae) on invertebrates and fish. - *River Res. Applics* 25: 734-744.
- Bilton, D. T., McAbendroth, L. C., Nicolet, P., Bedford, A., Rundle, S. D., Foggo, A. and Ramsay, P. M. 2009a. Ecology and conservation status of temporary and fluctuating ponds in two areas of southern England. - *Aquat. Conserv. mar. Freshwat. Ecosyst.* 19: 134-146.
- Biswas, J. K., Rana, S., Bhakta, J. N. and Jana, B. B. 2009a. Bioturbation potential of chironomid larvae for the sediment-water phosphorus exchange in simulated pond systems of varied nutrient enrichment. - *Ecol. Engng* 35: 1444-1453.
- Bitušík, P., Kubovčík, V., Štefková, E., Appleby, P. G. and Svitok, M. 2009a. Subfossil diatoms and chironomids along an altitudinal gradient in the High Tatra Mountain lakes: a multi-proxy record of past environmental trends. - *Hydrobiologia* 631: 65-85.
- Björk, P., Jin, S., Zhao, J., Singh, O. P., Persson, J.-O., Hellman, U. and Wieslander, L. 2009a. Specific combinations of SR proteins associate with single pre-messenger RNAs in vivo and contribute different functions. - *J. Cell Biol.* 184: 555-568.
- Blocksom, K. A. and Johnson, B. R. 2009a. Development of a regional macroinvertebrate index for large river bioassessment. - *Ecol. Indicators* 9: 313-328.
- Boakye, D. A., Fokam, E., Ghansah, A., Amakye, J., Wilson, M. D. and Brown, C. A. 2009a. *Cardiocladius oliffi* (Diptera: Chironomidae) as a potential biological control agent against *Simulium squamosum* (Diptera: Simuliidae). - *Parasites Vectors* 2, 20: 5 pp.
- Bogut, I., Vidakovic, J., Cerba, D. and Palijan, G. 2009a. Epiphytic meiofauna in stands of different submerged macrophytes. - *Ekoloji* 18, 70: 1-9.
- Boonsoong, B., Sangpradub, N. and Barbour, M. T. 2009a. Development of rapid bioassessment approaches using benthic macroinvertebrates for Thai streams. - *Envir. Monit. Assess.* 155: 129-147.
- Borza, P., Erős, T. and Oertel, N. 2009a. Food resource partitioning between two invasive gobiid species (Pisces, Gobiidae) in the littoral zone of the River Danube, Hungary. - *Int. Rev. Hydrobiol.* 94: 609-621.

- Bouchard Jr., R. W. and Ferrington Jr., L. C. 2009a. Winter growth, development, and emergence of *Diamesa mendotae* (Diptera: Chironomidae) in Minnesota streams. - *Envir: Ent.* 38: 250-259.
- Bouchard Jr., R. W. and Ferrington Jr., L. C. 2009b. Determination of Chironomidae thermal preferences and thermal partitioning among closely related taxa in a Minnesota stream using surface floating pupal exuviae. - In: Hughes, S. J., Khadem, M. and Carvalho, M. A. (eds.): *Proc. 16th Int. Chironomid Symp., Bolm Mus. munic. Funchal (Hist. nat.), Suppl. 13*: 191-198.
- Bredenhand, E. and Samways, M. J. 2009a. Impact of a dam on benthic macroinvertebrates in a small river in a biodiversity hotspot: Cape Floristic Region, South Africa. - *J. Insect Conserv.* 13: 297-307.
- Brodin, Y. and Andersson, M. H. 2009a. The marine splash midge *Telmatogon* [sic!] *japonicus* (Diptera; Chironomidae) - extreme and alien? - *Biol. Invasions* 11: 1311-1317.
- Buczko, K., Magyari, E. K., Bitušik, P. and Wacnik, A. 2009a. Review of dated Late Quaternary palaeolimnological records in the Carpathian Region, east-central Europe. - *Hydrobiologia* 631: 3-28.
- Bunbury, J. and Gajewski, K. 2009a. Postglacial climates inferred from a lake at treeline, southwest Yukon Territory, Canada. - *Quatern. Sci. Rev.* 28: 354-369.
- Burdett, A. S. and Watts, R. J. 2009a. Modifying living space: an experimental study of the influences of vegetation on aquatic invertebrate community structure - *Hydrobiologia* 618: 161-173.
- Burton, T. M. and Uzarski, D. G. 2009a. Biodiversity in protected coastal wetlands along the west coast of Lake Huron. - *Aquat. Ecosyst. Hlth Mgmt* 12: 63-76.
- Caldwell, B. A. 2009a. *Chironomus oceanicus* Packard, 1869, a new junior synonym of *Halocladius variabilis* (Staeger, 1839) (Diptera: Chironomidae). - In: Hughes, S. J., Khadem, M. and Carvalho, M. A. (eds.): *Proc. 16th Int. Chironomid Symp., Bolm Mus. munic. Funchal (Hist. nat.), Suppl. 13*: 207-212.
- Campbell, B. D., Haro, R. J. and Richardson, W. B. 2009a. Effects of agricultural land use on chironomid communities: comparisons among natural wetlands and farm ponds. - *Wetlands* 29: 1070-1080.
- Cañedo-Argüelles, M. 2009a. *Ecology of macroinvertebrate communities in transitional waters: Influence of the environment, response to disturbance and successional processes.* - Ph. D. Thes., Univ. Barcelona. 226 pp.
- Cañedo-Argüelles, M. and Rieradevall, M. 2009a. Quantification of environment-driven changes in epiphytic macroinvertebrate communities associated with *Phragmites australis*. - *J. Limnol.* 68: 229-241.
- Casatti, L., Júnior, J. L. V. e Ferreira, C. P. 2009a. Dieta do cascudo *Aspidoras fuscoguttatus* (Ostariophysi, Callichthyidae) em riachos com diferentes características limnológicas e estruturais. - *Biota neotrop.* 9, 1, bn02309012009: 113-122.
- Chan, E. K. W., Zhang, Y. and Dudgeon, D. 2009a. Substrate availability may be more important than aquatic insect abundance in the distribution of riparian orb-web spiders in the tropics. - *Biotropica* 41: 196-201.
- Charman, D. J., Barber, K. E., Blaauw, M., Langdon, P. G., Mauquoy, D., Daley, T. J., Hughes, P. D. M. and Karofeld, E. 2009a. Climate drivers for peatland palaeoclimate records. - *Quatern. Sci. Rev.* 28: 1811-1819.
- Cheng, M. and Wang, X. 2009a. *Thienemannimyia* Fittkau (Diptera: Chironomidae) from China. - *Zootaxa* 2074: 50-60.
- Chibunda, R. T. 2009a. Chronic toxicity of mercury (HgCl₂) to the benthic midge *Chironomus riparius*. - *Int. J. envir. Res.* 3: 455-462.
- Choi, J. and Ha, M.-H. 2009a. Effect of cadmium exposure on the globin protein expression in 4th instar larvae of *Chironomus riparius* Mg. (Diptera: Chironomidae): An ecotoxicoproteomics approach. - *Proteomics* 9: 31-39.
- Chowdhury, N., Chatterjee, S. K., Laskar, S. and Chandra, G. 2009a. Larvicidal activity of *Solanum villosum* Mill (Solanaceae: Solanales) leaves to *Anopheles subpictus* Grassi (Diptera: Culicidae) with effect on non-target *Chironomus circumdatus* Kieffer (Diptera: Chironomidae). - *J. Pest Sci.* 82: 13-18.
- Como, S. and Magni, P. 2009a. Temporal changes of a macrobenthic assemblage in

- harsh lagoon sediments. - *Estuar. coast. Shelf Sci.* 83: 638-646.
- Corbi, J. J. and Trivinho-Strixino, S. 2009a. Effects of land use on lotic chironomid communities of southeast of Brazil: emphasis on the impact of sugar cane cultivation. - In: Hughes, S. J., Khadem, M. and Carvalho, M. A. (eds.): *Proc. 16th Int. Chironomid Symp., Bolm Mus. munic. Funchal (Hist. nat.), Suppl. 13*: 93-100.
- Corcoran, R. M., Lovvorn, J. R. and Heglund, P. J. 2009a. Long-term change in limnology and invertebrates in Alaskan boreal wetlands. - *Hydrobiologia* 620: 77-89.
- Cranston, P. S. 2009a. A new genus of trans-Tasman midge: *Anzacladius* gen. n. (Diptera: Chironomidae: Orthocladiinae). - *Aust. J. Ent.* 48: 130-139.
- Cuppen, H., Klink, A. and Moller Pillot, H. 2009a. The larvae of *Graceus ambiguus* and *Sergentia near prima* and their identification. - *Lauterbornia* 67: 29-37.
- Datkhile, K. D., Dongre, T. K., Mukhopadhyaya, R. and Nath, B. B. 2009a. Gamma radiation tolerance of a tropical species of midge, *Chironomus ramosus* Chaudhuri (Diptera: Chironomidae). - *Int. J. Radiat. Biol.* 85: 495-503.
- Datkhile, K. D., Mukhopadhyaya, R., Dongre, T. K. and Nath, B. B. 2009a. Increased level of superoxide dismutase (SOD) activity in larvae of *Chironomus ramosus* (Diptera: Chironomidae) subjected to ionizing radiation. - *Comp. Biochem. Physiol. Pt C Toxic. Pharmacol.* 149: 500-506.
- De Jonge, M., Dreesen, F., De Paepe, J., Blust, R. and Bervoets, L. 2009a. Do acid volatile sulfides (AVS) influence the accumulation of sediment-bound metals to benthic invertebrates under natural field conditions? - *Envir. Sci. Technol.* 43: 4510-4516.
- De Lange, H. J., Peeters, E. T. H. M., Harmsen, J., Maas, H. and De Jonge, J. 2009a. Seasonal variation of total and biochemically available concentrations of PAHs in a floodplain lake sediment has no effect on the benthic invertebrate community. - *Chemosphere* 75: 319-326.
- Ding, Y., You, J. and Lydy, M. J. 2009a. Analysis of pyrethroid insecticides in *Chironomus dilutus* using matrix solid phase dispersion extraction. - *Bull. envir. Contam. Toxic.* 83: 388-392.
- Di Veroli, A., Goretti, E., Marcucci, C., Fabrizi, A., Scopetta, L. and Di Giovanni, M. V. 2009a. Incidence of larvae mouthpart deformities in *Chironomus plumosus* (DIPTERA: CHIRONOMIDAE) and *Procladius* sp. (DIPTERA: CHIRONOMIDAE) from Piediluco Lake, Italy. - In: Hughes, S. J., Khadem, M. and Carvalho, M. A. (eds.): *Proc. 16th Int. Chironomid Symp., Bolm Mus. munic. Funchal (Hist. nat.), Suppl. 13*: 13-20.
- Dörner, H., Skov, C., Berg, S., Schulze, T., Beare, D. J. and van der Velde, G. 2009a. Piscivory and trophic position of *Anguilla anguilla* in two lakes: importance of macrozoobenthos density. - *J. Fish Biol.* 74: 2115-2131.
- Dombrovskii, K. O. 2009a. Znachenie dvustvorchatykh mollyuskov v obrazovanii konsortsii vodnykh bespozvonochnykh v litorali iskusstvennogo evtrofnogo ozero. (Role of bivalve mollusks in formation of aquatic invertebrate consortia in the littoral zone of an artificial lake.) - *Ekologiya* 2009, 2: 127-132. [also published in: *Russ. J. Ecol.* 40: 116-120.]
- Domingues, I., Guilhermino, L., Soares, A. M. V. M., Nogueira, A. J. A. and Monaghan, K. A. 2009a. Influence of exposure scenario on pesticide toxicity in the midge *Kiefferulus calligaster* (Kieffer). - *Ecotoxic. envir. Saf.* 72: 450-457.
- Dornfeld, C. B., Moreira-Santos, M., Espíndola, E. L. G. and Ribeiro, R. 2009a. Do larvae and ovipositing *Chironomus riparius* (Diptera: Chironomidae) females avoid copper-contaminated environments? - *Hum. ecol. Risk Assess.* 15: 63-75.
- Dubois, M. and Hare, L. 2009a. Selenium assimilation and loss by an insect predator and its relationship to Se subcellular partitioning in two prey types. - *Envir. Pollut.* 157: 772-777.
- Dubois, M. and Hare, L. 2009b. Subcellular distribution of cadmium in two aquatic invertebrates: change over time and relationship to Cd assimilation and loss by a predatory insect. - *Envir. Sci. Technol.* 43: 356-361.
- Dukowska, M., Grzybkowska, M., Marszał, L. and Zięba, G. 2009a. The food preferences of three-spined stickleback, *Gasterosteus aculeatus* L., downstream from a dam

- reservoir. - *Oceanol. hzydrobiol. Stud.* 38: 39-50.
- Durance, I. and Ormerod, S. J. 2009a. Trends in water quality and discharge confound long-term warming effects on river macroinvertebrates. - *Freshwat. Biol.* 54: 388-405.
- Durnova, N. A. 2009a. Khromosomnyi polimorfizm i tsitotipy *Endochironomus tendens* F. (Diptera, Chironomidae) iz vodoemov Saratovskoi i Samarskoi oblasti. (Chromosome polymorphism and cytotypes of *Endochironomus tendens* F. (Diptera, Chironomidae) from Saratov and Samara regions.) - *Tsitologiya* 51: 592-601. [also published in: *Cell Tissue Biol.* 3: 340-348.]
- Dussault, E. B., Balakrishnan, V. K., Borgmann, U., Solomon, K. R. and Sibley, P. K. 2009a. Bioaccumulation of the synthetic hormone 17 α -ethinylestradiol in the benthic invertebrates *Chironomus tentans* and *Hyalella azteca*. - *Ecotoxic. envir. Saf.* 72: 1635-1641.
- Dussault, E. B., Balakrishnan, V. K., Solomon, K. R. and Sibley, P. K. 2009a. Matrix effects on mass spectrometric determinations of four pharmaceuticals and personal care products in water, sediments, and biota. - *Can. J. Chem.* 87: 662-672.
- Ekrem, T. and Stur, E. 2009a. A review of the genus *Zavrelia* (Diptera: Chironomidae). - *Eur. J. Ent.* 106: 119-144.
- Eldridge, J. L., Krapu, G. L. and Johnson, D. H. 2009a. Feeding ecology of Arctic-nesting sandpipers during spring migration Through the Prairie Pothole Region. - *J. Wildl. Mgmt* 73: 248-252.
- Engels, S., Bohncke, S. J. P., Bos, J. A. A., Helmens, K. and Heiri, O. 2009a. Chironomid-based inferences of local and regional environmental change during the early Middle Weichselian in Northeast Finland. - In: Hughes, S. J., Khadem, M. and Carvalho, M. A. (eds.): *Proc. 16th Int. Chironomid Symp., Bolm Mus. munic. Funchal (Hist. nat.), Suppl. 13*: 185-190.
- Ermolaeva, O. V. 2009a. Kariotip *Diamesa hamaticornis* Kieffer, 1924 (Diptera, Chironomidae) i sravnitel'nyi kariologicheskii analiz vidov v rode *Diamesa* Meigen. (The karyotype of *Diamesa hamaticornis* Kieffer, 1924 (Diptera, Chironomidae) and a comparative karyological analysis of *Diamesa* Meigen species. - *Evraziat. ent. Zh. (= Euroas. ent. J.)* 8, Suppl. 1: 29-32.
- Fernández, H. R., Romero, F. and Domínguez, E. 2009a. Intermountain basins use in subtropical regions and their influences on benthic fauna. - *River Res. Applics* 25: 181-193.
- Ferns, P. N. and Reed, J. P. 2009a. Effects of the Cardiff Bay tidal barrage on the abundance, ecology and behaviour of shelducks *Tadorna tadorna*. - *Aquat. Conserv. mar. Freshwat. Ecosyst.* 19: 466-473.
- Ferrington Jr., L. C., Masteller, E. C. and Santiago-Blay, J. A. 2009a. Comparison and significance of Chironomidae emergence from Lake Erie and Presque Isle Bay, Erie, Pennsylvania, U. S. A. - In: Hughes, S. J., Khadem, M. and Carvalho, M. A. (eds.): *Proc. 16th Int. Chironomid Symp., Bolm Mus. munic. Funchal (Hist. nat.), Suppl. 13*: 21-28.
- Filippov, A. A. and Riedel, F. 2009a. The late Holocene mollusc fauna of the Aral Sea and its biogeographical and ecological interpretation. - *Limnologica* 39: 67-85.
- Free, G., Bowman, J., McGarrigle, M., Caroni, R., Donnelly, K., Tierney, D., Trodd, W. and Little, R. 2009a. The identification, characterization and conservation value of isoetid lakes in Ireland. - *Aquat. Conserv. mar. Freshwat. Ecosyst.* 19: 264-273.
- Free, G., Solimini, A. G., Rossaro, B., Marziali, L., Giacchini, R., Paracchini, B., Ghiani, M., Vaccaro, S., Gawlik, B. M., Fresner, R., Santner, G., Schönhuber, M. and Cardoso, A. C. 2009a. Modelling lake macroinvertebrate species in the shallow sublittoral: relative roles of habitat, lake morphology, aquatic chemistry and sediment composition. - *Hydrobiologia* 633: 123-136.
- French III, J. R. P., Schaeffer, J. S., Roseman, E. F., Kiley, C. S. and Fouilleroux, A. 2009a. Abundance and distribution of benthic macroinvertebrates in offshore soft sediments in Western Lake Huron, 2001-2007. - *J. Gt Lakes Res.* 35: 120-127.
- Friberg, N., Dybkjær, J. B., Olafsson, J. S., Gislason, G. M., Larsen, S. E. and Lauridsen, T. L. 2009a. Relationships between structure and function in streams contrasting in temperature. - *Freshwat. Biol.* 54: 2051-2068.
- Frouz, J., Lobinske, R. J. and Ali, A. 2009a. Intraspecific competition of *Glyptotendipes paripes* (Diptera: Chironomidae) under

- laboratory conditions. - *Aquat. Ecol.* 43: 487-500.
- Fu, Y. and Wang, X. 2009a. Four new species of *Nanocladius* Kieffer from Oriental China (Diptera: Chironomidae: Orthoclaadiinae). - *Zootaxa* 1985: 43-51.
- Fusari, L. M., Roque, F. O. and Hamada, N. 2009a. *Oukuriella pesae*, new species of sponge-dwelling chironomid (Insecta: Diptera) from Amazonia, Brazil. - *Zootaxa* 2146: 61-68.
- Gallardo, B., Gascón, S., García, M. and Comín, F. A. 2009a. Testing the response of macroinvertebrate functional structure and biodiversity to flooding and confinement. - *J. Limnol.* 68: 315-326.
- Garbary, D. J., Jamieson, M. M. and Taylor, B. R. 2009a. Population ecology of the marine insect *Halocladus variabilis* (Diptera: Chironomidae) in the rocky intertidal zone of Nova Scotia, Canada. - *Mar. Ecol. Prog. Ser.* 376: 193-202.
- Gathorne-Hardy, F. J., Erlendsson, E., Langdon, P. G. and Edwards, K. J. 2009a. Lake sediment evidence for late Holocene climate change and landscape erosion in western Iceland. - *J. Paleolimnol.* 42: 413-426.
- Giddings, J. M., Barber, I. and Warren-Hicks, W. 2009a. Comparative aquatic toxicity of the pyrethroid insecticide lambda-cyhalothrin and its resolved isomere gamma-cyhalothrin. - *Ecotoxicology* 18: 239-249.
- Gilka, W. 2009a. Order Diptera, family Chironomidae - tribe Tanytarsini. - In: Harten, A. van (ed.): *Arthropod fauna of the United Arab Emirates, vol. 2*: 667-682. Dar Al Ummah Print., Publ., Distrib., Advertis., Abu Dhabi.
- Gilka, W. and Paasivirta, L. 2009a. Evaluation of diagnostic characters of the *Tanytarsus chinyensis* group (Diptera: Chironomidae), with description of a new species from Lapland. - *Zootaxa* 2197: 31-42.
- Gillies, C. L., Hose, G. C. and Turak, E. 2009a. What do qualitative rapid assessment collections of macroinvertebrates represent? A comparison with extensive quantitative sampling. - *Envir. Monit. Assess.* 149: 99-112.
- Gonçalves, A. L. and Canhoto, C. 2009a. Decomposition of eucalypt and alder mixtures: responses to variation in evenness. - *Fundam. appl. Limnol.* 173: 293-303.
- Grabowska, J., Grabowski, M. and Kostecka, A. 2009a. Diet and feeding habits of monkey goby (*Neogobius fluviatilis*) in a newly invaded area. - *Biol. Invasions* 11: 2161-2170.
- Grabowska, J., Grabowski, M., Pietraszewski, D. and Gmur, J. 2009a. Non-selective predator - the versatile diet of Amur sleeper (*Percottus glenii* Dybowski, 1877) in the Vistula River (Poland), a newly invaded ecosystem. - *J. appl. Ichthyol.* 25: 451-459.
- Green, P. 2009a. The response of chironomids to water and sediment chemistry in urban canals. - In: Hughes, S. J., Khadem, M. and Carvalho, M. A. (eds.): *Proc. 16th Int. Chironomid Symp., Bolm Mus. munic. Funchal (Hist. nat.), Suppl. 13*: 29-37.
- Gremyatchikh, V. A., Tomilina, I. I. i Grebenyuk, L. P. 2009a. Vliyanie khlorida rtuti na morfofunksional'nye pokazateli lichinok *Chironomus riparius* Meigen (Diptera, Chironomidae). (The effect of mercury chloride on morphofunctional parameters in *Chironomus riparius* Meigen (Diptera, Chironomidae) larvae.) - *Biol. vnutr. Vod* 2009, 1: 94-101. [also published in: *Inland Wat. Biol.* 2,1: 89-95.]
- Gresens, S. E. and Ferrington Jr., L. C. 2009a. Taxonomic distinctness provides an alternative view of the diversity of Chironomidae (Diptera) assemblages. - In: Hughes, S. J., Khadem, M. and Carvalho, M. A. (eds.): *Proc. 16th Int. Chironomid Symp., Bolm Mus. munic. Funchal (Hist. nat.), Suppl. 13*: 101-108.
- Grohs, K. L., Klumb, R. A., Chipps, S. R. and Wanner, G. A. 2009a. Ontogenetic patterns in prey use by pallid sturgeon in the Missouri River, South Dakota and Nebraska. - *J. appl. Ichthyol.* 25, s2: 48-53.
- Guilizzoni, P., Massafiero, J., Lami, A., Piovano, E. L., Guevara, S. R., Formica, S. M., Daga, R., Rizzo, A. and Gerli, S. 2009a. Palaeolimnology of Lake Hess (Patagonia, Argentina): multi-proxy analyses of short sediment cores. - *Hydrobiologia* 631: 289-312.
- Gunderina, L. I., Kiknadze, I. I., Istomina, A. G. and Butler, M. 2009a. Geograficheskaya differentsiatsiya genomnoi DNK *Chironomus plumosus* (Diptera, Chironomidae) v prirodnykh populyatsiyakh Golarktiki. (Geographic differentiation of genomic DNA of *Chironomus plumosus* (Diptera, Chironomidae) in natural holarctic

- populations.) - *Genetika* 45: 64-72. [also published in: *Russ. J. Genet.* 45: 54-62.]
- Gunkel, G., Beulker, C., Hoffmann, A. and Kosmol, J. 2009a. Fine particulate organic matter (FPOM) transport and processing in littoral interstices - use of fluorescent markers. - *Limnologica* 39: 185-199.
- Halpern, M., Shakéd, T., Pukall, R. and Schumann, P. 2009a. *Leucobacter chironomi* sp. nov., a chromate-resistant bacterium isolated from a chironomid egg mass. - *Int. J. syst. evolut. Microbiol.* 50: 665-670.
- Hanson, M. A., Bowe, S. E., Ossman, F. G., Fieberg, J., Butler, M. G. and Koch, R. 2009a. Influences of forest harvest and environmental gradients on aquatic invertebrate communities of seasonal ponds. - *Wetlands* 29: 884-895.
- Hargrave, C. W. 2009a. Effects of fish species richness and assemblage composition on stream ecosystem function. - *Ecol. Freshwat. Fish* 18: 24-32.
- Hargrave, C. W., Gary, K. P. and Rosado, S. K. 2009a. Potential effects of elevated atmospheric carbon dioxide on benthic autotrophs and consumers in stream ecosystems: a test using experimental stream mesocosms. - *Global Change Biol.* 15: 2779-2790.
- Harwood, A. D., You, J. and Lydy, M. J. 2009a. Temperature as a toxicity identification evaluation tool for pyrethroids insecticides: toxicokinetic confirmation. - *Envir. Toxic. Chem.* 28: 1051-1058.
- Heino, J., Ilmonen, J., Kotanen, J., Mykrä, H., Paasivirta, L., Soininen, J. and Virtanen, R. 2009a. Surveying biodiversity in protected and managed streams: Algae, macrophytes and macroinvertebrates in boreal forest streams. - *Ecol. Indicators* 9: 1179-1187.
- Heiri, O. and Lotter, A. F. 2009a. Chironomidae (Diptera) in Alpine lakes: a study of subfossil assemblages in lake surface sediments. - In: Hughes, S. J., Khadem, M. and Carvalho, M. A. (eds.): *Proc. 16th Int. Chironomid Symp., Bolm Mus. munic. Funchal (Hist. nat.), Suppl. 13: 177-184.*
- Helmens, K. F., Risberg, J., Jansson, K. N., Weckström, J., Berntsson, A., Tillman, P. K., Johansson, P. W. and Wastegård, S. 2009a. Early MIS 3 glacial lake evolution, ice-marginal retreat pattern and climate at Sokli (northeastern Fennoscandia). - *Quatern. Sci. Rev.* 28: 1880-1894.
- Hessle, V., Björk, P., Sokolowski, M., Valdivia, E. G. de, Silverstein, R., Artemenko, K., Tyagi, A., Maddalo, G., Ilag, L., Helbig, R., Zubarev, R. A. and Visa, N. 2009a. The exosome associates cotranscriptionally with the nascent pre-mRNP through interactions with heterogeneous nuclear ribonucleoproteins. - *Molec. Biol. Cell* 20: 3459-3470.
- Henriques-Oliveira, A. L., Silva, R. A. and Nessimian, J. L. 2009a. First recorded [sic!] of *Pontomyia* Edwards, 1926 (Diptera: Chironomidae: Tanytarsini) in Brazil. - *Biota neotrop.* 9,1, bn02309012009: 271-274.
- Hirabayashi, K., Yoshizawa, K., Oga, K., Yoshida, N., Ariizumi, K. and Kazama, F. 2009a. Change of chironomid fauna (Diptera: Chironomidae) in eutrophic Lake Kawaguchi, Japan. - In: Hughes, S. J., Khadem, M. and Carvalho, M. A. (eds.): *Proc. 16th Int. Chironomid Symp., Bolm Mus. munic. Funchal (Hist. nat.), Suppl. 13: 109-117.*
- Hoang, T. C., Schuler, L. J., Rogevich, E. C., Bachman, P. M., Rand, G. M. and Frakes, R. A. 2009a. Copper release, speciation, and toxicity following multiple floodings of copper enriched agriculture soils: implications in Everglades restoration. - *Wat. Air Soil Pollut.* 199: 79-93.
- Höckelmann, C., Becher, P. G., Von Reuß, S. H. and Jüttner, F. 2009a. Sesquiterpenes of the geosmin-producing cyanobacterium *Calothrix* PCC 7507 and their toxicity to invertebrates. - *Z. Naturforsch. Sect. C, J. Biosci.* 64: 49-55.
- Hoekman, D., Winston, R. and Mitchell, N. 2009a. Top-down and bottom-up effects of a processing detritivore. - *J. N. Am. benthol. Soc.* 28:552-559.
- Holmes, N., Langdon, P. G. and Caseldine, C. J. 2009a. Subfossil chironomid variability in surface sediment samples from Icelandic lakes: implications for the development and use of training sets. - *J. Paleolimnol.* 42: 281-295.
- Horikawa, D. D., Iwata, K. I., Kawai, K., Koseki, S., Okuda, T. and Yamamoto, K. 2009a. High hydrostatic pressure tolerance of four different anhydrobiotic animal species. - *Zool. Sci.* 26: 238-242.
- Hormes, A., Blaauw, M., Dahl, S. O., Nesje, A. and Possnert, G. 2009a. Radiocarbon wiggle-match dating of proglacial lake sediments - Implications for the 8.2 ka event. - *Quatern. Geochronol.* 4: 267-277.

- Hu, Z.-J., Liu, Q.-G., Chen, L.-J. and Peng, Z.-R. 2009a. (Structural characteristics of chironomid community and their indicative significance in bioassessment of water quality in Mingzhu Lake of Chongming Island, Shanghai.) - *Chin. J. appl. Ecol.* 20: 929-936.
- Huddleston III, G. M., Dorn, P. B., Gillespie Jr., W. B., Wong, D. C. and Slocomb, J. P. 2009a. Assessment of the ecological effects of arsenic on a southern Ohio, USA stream. - *Integr. envir. Assess. Mgmt* 5: 302-319.
- Hughes, S. J., Khadem, M. and Carvalho, M. A. 2009a. Foreword. - In: Hughes, S. J., Khadem, M. and Carvalho, M. A. (eds.): *Proc. 16th Int. Chironomid Symp., Bolm Mus. munic. Funchal (Hist. nat.), Suppl. 13*: 1.
- Hunter, W., Yang, Y., Reichenberg, F., Mayer, P. and Gan, J. 2009a. Measuring pyrethroids in sediment porewater using matrix-solid phase microextraction. - *Envir. Toxic. Chem.* 28: 36-43.
- Hutchens Jr., J. J., Schuldt, J. A., Richards, C., Johnson, L. B., Host, G. E. and Breneman, D. H. 2009a. Multi-scale mechanistic indicators of Midwestern USA stream macroinvertebrates. - *Ecol. Indicators* 9: 1138-1150.
- Iimonen, J., Paasivirta, L., Virtanen, R. and Muotka, T. 2009a. Regional and local drivers of macroinvertebrate assemblages in boreal springs. - *J. Biogeogr.* 36: 822-834.
- Ilyashuk, B., Gobet, E., Heiri, O., Lotter, A. F., van Leeuwen, J. F. N., van der Knaap, W. O., Ilyashuk, E., Oberli, F. and Ammann, B. 2009a. Lateglacial environmental and climatic changes at the Maloja Pass, Central Swiss Alps, as recorded by chironomids and pollen. - *Quatern. Sci. Rev.* 28: 1340-1353.
- Inoue, E., Kimura, G., Denda, M., Tokioka, T., Tsushima, K., Amano, K. and Hirabayashi, K. 2009a. Usefulness and problems of larval rearing for assessment of lotic chironomid assemblages: a case study in a riffle/pool section at the middle reaches of the Shinano River, Japan. - In: Hughes, S. J., Khadem, M. and Carvalho, M. A. (eds.): *Proc. 16th Int. Chironomid Symp., Bolm Mus. munic. Funchal (Hist. nat.), Suppl. 13*: 119-126.
- Ioannou, C. C. and Krause, J. 2009a. Interactions between background matching and motion during visual detection can explain why cryptic animals keep still. - *Biol. Lett.* 5: 191-193.
- Istomina, A. G., Kiknadze, I. I. i Gunderina, L. I. 2009a. Khromosomnaya izmenchivost' v prirodnykh populyatsiyakh *Chironomus cingulatus* Meigen (Diptera, Chironomidae). (Chromosomal variability in natural populations of *Chironomus cingulatus* Meigen (Diptera, Chironomidae).) - *Tsitologiya* 51: 138-148. [also published in: *Cell Tissue Biol.* 3: 149-161.]
- Jabłońska-Barna, I. and Michailova, P. 2009a. Cytogenetic characteristics of species of the *Chironomus* genus (Diptera, Chironomidae) from Lake Łuknajno Biosphere Reserve (northwest Poland). - *Oceanol. hydrobiol. Stud.* 38: 77-81.
- Jacobsen, R. E. 2009a. Midge (DIPTERA: CHIRONOMIDAE and CERATOPOGONIDAE) community response to canal discharge into Everglades National Park, Florida. - In: Hughes, S. J., Khadem, M. and Carvalho, M. A. (eds.): *Proc. 16th Int. Chironomid Symp., Bolm Mus. munic. Funchal (Hist. nat.), Suppl. 13*: 39-50.
- James, A. B. W., Dewson, Z. S. and Death, R. G. 2009a. The influence of flow reduction on macroinvertebrate drift density and distance in three New Zealand streams. - *J. N. Am. benthol. Soc.* 28: 220-232.
- James, R. 2009a. Temporal costs of feeding and predation rates in *Betta splendens* (Regan) in relation to body weight, feed type and sex. - *Trop. Life Sci. Res.* 20: 7-16.
- Jungmann, D., Bandow, C., Gildemeister, T., Nagel, R., Preuss, T. G., Ratte, H. T., Shinn, C., Weltje, L. and Maes, H. M. 2009a. Chronic toxicity of fenoxycarb to the midge *Chironomus riparius* after exposure in sediments of different composition. - *J. Soils Sediments* 9: 94-102.
- Kaiser, T. S. and Heckel, D. G. 2009a. Isolation of 10 polymorphic microsatellite loci in the marine midge *Clunio marinus* (Chironomidae, Diptera) and their efficient characterization by heteroduplex analysis. - *Molec. Ecol. Resour.* 9: 195-198.
- Karouna-Renier, N. K. and Rao, K. R. 2009a. An inducible HSP70 gene from the midge *Chironomus dilutus*: characterization and transcription profile under environmental stress. - *Insect molec. Biol.* 18: 87-96.
- Kiknadze, I. I. I Broshkov, A. D. 2009a. Kariotip i khromosomnyi polimorfizm u khironomid *Chironomus melanescens* Keyl, 1961 (Diptera, Chironomidae). - (The karyotype

- and chromosomal polymorphism of the midge *Chironomus melanescens* Keyl, 1961 (Diptera, Chironomidae). - *Evrasiat. ent. Zh.* (= *Euroas. ent. J.*) 8, *Suppl. 1*: 19-28.
- Kirilova, E., Heiri, O., Enters, D., Cremer, H., Lotter, A. F., Zolitschka, B. and Hübener, T. 2009a. Climate-induced changes in the trophic status of a Central European lake. - *J. Limnol.* 68: 71-82.
- Klug, M., Schmidt, S., Bennike, O., Heiri, O., Melles, M. and Wagner, B. 2009a. Lake sediments from Store Koldewey, Northeast Greenland, as archive of Late Pleistocene and Holocene climatic and environmental changes. - *Boreas* 38: 59-71.
- Koodalingam, A., Mullainadhan, P. and Arumugam, M. 2009a. Antimosquito activity of aqueous kernel extract of soapnut *Sapindus emarginatus*: impact on various developmental stages of three vector mosquito species and nontarget aquatic insects. - *Parasit. Res.* 105: 1425-1434.
- Koperski, P. 2009a. Reduced diversity and stability of chironomid assemblages (Chironomidae, Diptera) as the effect of moderate stream degradation. - *Pol. J. Ecol.* 57: 125-138.
- Koszalka, J. 2009a. Chironomidae (Diptera) community response to improved oxygen conditions in a naturally recovering lake (Łękuk Wielki, northeast Poland). - *Oceanol. hydrobiol. Stud.* 38: 83-89.
- Krashennikov, A. B. i Makarchenko, E. A. 2009a. Novye i maloizvestnyye vidy khironomid (Diptera, Chironomidae, Orthoclaadiinae) iz Visherskogo gosudarstvennogo prirodnogo zapovednika (Severnyi Ural). (New and little-known species of chironomids (Diptera, Chironomidae, Orthoclaadiinae) of the Vishersky Nature Reserve (North Ural).) - *Evrasiat. ent. Zh.* (= *Euroas. ent. J.*) 8, *Suppl. 1*: 97-101.
- Krosch, M. N., Baker, A. M., McKie, B. G., Mather, P. B. and Cranston, P. S. 2009a. Deeply divergent mitochondrial lineages reveal patterns of local endemism in chironomids of the Australian Wet Tropics. - *Austral Ecol.* 34: 317-328.
- Kurek, J. and Cwynar, L. C. 2009a. Effects of within-lake gradients on the distribution of fossil chironomids from maar lakes in western Alaska: implications for environmental reconstructions. - *Hydrobiologia* 623: 37-52.
- Kurek, J. and Cwynar, L. C. 2009b. The potential of site-specific and local chironomid-based inference models for reconstructing past lake levels. - *J. Paleolimnol.* 42: 37-50.
- Kurek, J., Cwynar, L. C., Ager, T. A., Abbott, M. B. and Edwards, M. E. 2009a. Late Quaternary paleoclimate of western Alaska inferred from fossil chironomids and its relation to vegetation histories. - *Quatern. Sci. Rev.* 28: 799-811.
- Kurek, J., Cwynar, L. C. and Vermaire, J. C. 2009a. A late Quaternary paleotemperature record from Hanging Lake, northern Yukon Territory, eastern Beringia. - *Quatern. Res.* 72: 246-257.
- Kwan, J. A., Novak, M. G., Hyles, T. S. and Niemela, M. K. 2009a. Mortality of nontarget arthropods from an aerial application of pyrethrins. - *J. Am. Mosquito Control Ass.* 25: 218-220.
- Lagauzère, S., Boyer, P., Stora, G. and Bonzom, J.-M. 2009a. Effects of uranium-contaminated sediments on the bioturbation activity of *Chironomus riparius* larvae (Insecta, Diptera) and *Tubifex tubifex* worms (Annelida, Tubificidae). - *Chemosphere* 76: 324-334.
- Lagauzère, S., Pischedda, L., Cuny, P., Gilbert, F., Stora, G. and Bonzom, J.-M. 2009a. Influence of *Chironomus riparius* (Diptera, Chironomidae) and *Tubifex tubifex* (Annelida, Oligochaeta) on oxygen uptake by sediments. Consequences of uranium contamination. - *Envir. Pollut.* 157: 1234-1242.
- Lamentowicz, M., Balwierz, Z., Forsytek, J., Płóciennik, M., Kittel, P., Kloss, M., Twardy, J., Zurek, S. and Pawlyta, J. 2009a. Multiproxy study of anthropogenic and climatic changes in the last two millennia from a small mire in central Poland. - *Hydrobiologia* 631: 213-230.
- Langford, T. E. L., Shaw, P. J., Ferguson, A. J. D. and Howard, S. R. 2009a. Long-term recovery of macroinvertebrate biota in grossly polluted streams: Re-colonisation as a constraint to ecological quality. - *Ecol. Indicators* 9: 1064-1077.
- Langton, P. H. 2009a. A structural hermaphrodite *Micropsectra* (DIPTERA: CHIRONOMIDAE). - In: Hughes, S. J., Khadem, M. and Carvalho, M. A. (eds.): *Proc. 16th Int. Chironomid Symp., Bolm Mus. munic. Funchal (Hist. nat.), Suppl. 13*: 213-216.

- Larocque, I., Grosjean, M., Heiri, O., Bigler, C. and Blass, A. 2009a. Comparison between chironomid-inferred July temperatures and meteorological data AD 1850–2001 from varved Lake Silvaplana, Switzerland. - *J. Paleolimnol.* 41: 329-342.
- Larsen, S., Vaughan, I. P. and Ormerod, S. J. 2009a. Scale-dependent effects of fine sediments on temperate headwater invertebrates. - *Freshwat. Biol.* 54: 203-219.
- Lau, D. C. P., Leung, K. M. Y. and Dudgeon, D. 2009a. What does stable isotope analysis reveal about trophic relationships and the relative importance of allochthonous and autochthonous resources in tropical streams? A synthetic study from Hong Kong. - *Freshwat. Biol.* 54: 127-141.
- Laufer, G., Arim, M., Loureiro, M., Piñeiro-Guerra, J.M., Clavijo-Baquet, S. and Fagúndez, C. 2009a. Diet of four annual killifishes: An intra and interspecific comparison. - *Neotrop. Ichthyol.* 7: 77-86.
- Lee, S.-W. and Choi, J. 2009a. Multi-level ecotoxicity assay on the aquatic midge, *Chironomus tentans* (Diptera, Chironomidae) exposed to octachlorostyrene. - *Envir. Toxic. Pharmac.* 28: 269-274.
- Lee, S.-W., Kim, S.-M. and Choi, J. 2009a. Genotoxicity and ecotoxicity assays using the freshwater crustacean *Daphnia magna* and the larva of the aquatic midge *Chironomus riparius* to screen the ecological risks of nanoparticle exposure. - *Envir. Toxic. Pharmac.* 28: 86-91.
- Lencioni, V., Boschini, D. and Rebecchi, L. 2009a. Expression of the 70 kDa Heat shock protein family in Alpine freshwater chironomids (Diptera, Chironomidae) under natural conditions. - *J. Limnol.* 68: 251-256.
- Lencioni, V., Marziali, L. and Rossaro, B. 2009a. Hyporheic chironomids in alpine streams. - In: Hughes, S. J., Khadem, M. and Carvalho, M. A. (eds.): *Proc. 16th Int. Chironomid Symp., Bolm Mus. munic. Funchal (Hist. nat.), Suppl. 13*: 127-132.
- León Paumen, M., de Voogt, P., van Gestel, C. A. M. and Kraak, M. H. S. 2009a. Comparative chronic toxicity of homo- and heterocyclic aromatic compounds to benthic and terrestrial invertebrates: Generalizations and exceptions. - *Sci. tot. Envir.* 407: 4605-4609.
- Leonard, E. M., Pierce, L. M., Gillis, P. L., Wood, C. M. and O'Donnell, M. J. 2009a. Cadmium transport by the gut and Malpighian tubules of *Chironomus riparius*. - *Aquat. Toxic.* 92: 179-186.
- Li, A. O. Y., Ng, L. C. Y. and Dudgeon, D. 2009a. Effects of leaf toughness and nitrogen content on litter breakdown and macroinvertebrates in a tropical stream. - *Aquat. Sci.* 71: 80-93.
- Lobo-Rodrigues, J. J. 2009a. *Taxonomia da fase adulta das espécies brasileiras de Goeldichironomus Fittkau, 1965 com redescritções de dez espécies e chave dicotômica (INSECTA: DIPTERA, CHIRONOMIDAE)*. - Tes. Mestr., Univ. Fed. Rio de Janeiro. 108 pp.
- Ludlam, J. P. and Magoulick, D. D. 2009a. Spatial and temporal variation in the effects of fish and crayfish on benthic communities during stream drying. - *J. N. Am. benthol. Soc.* 28: 371-382.
- Luoto, T. P. 2009a. Subfossil Chironomidae (Insecta: Diptera) along a latitudinal gradient in Finland: development of a new temperature inference model. - *J. Quatern. Sci.* 24: 150-158.
- Luoto, T. P. 2009b. A Finnish chironomid- and chaoborid-based inference model for reconstructing past lake levels. - *Quatern. Sci. Rev.* 28: 1481-1489.
- Maasri, A., Fayolle, S. and Franquet, E. 2009a. Epilithic chironomid community structure: is it a matter of algae? - In: Hughes, S. J., Khadem, M. and Carvalho, M. A. (eds.): *Proc. 16th Int. Chironomid Symp., Bolm Mus. munic. Funchal (Hist. nat.), Suppl. 13*: 133-140.
- MacDonald, G. M., Porinchu, D. F., Rolland, N., Kremenetsky, K. V. and Kaufman, D. S. 2009a. Paleolimnological evidence of the response of the central Canadian treeline zone to radiative forcing and hemispheric patterns of temperature change over the past 2000 years. - *J. Paleolimnol.* 41: 129-141.
- Majumdar, T. N. and Gupta, A. 2009a. Acute toxicity of endosulfan and malathion on *Chironomus ramosus* (Insecta: Diptera: Chironomidae) from north Cachar hills, Assam, India. - *J. envir. Biol.* 30: 469-470.
- Makarchenko, E. A. and Endo, K. 2009a. The description of immature stages of *Linevitshia* Makarchenko and *Sasayusurika* Makarchenko (Diptera, Chironomidae, Diamesinae), with some remarks on taxonomy and systematics

- of these genera. - *Evraziat. ent. Zh. (= Euroas. ent. J.)* 8, *Suppl. 1*: 64-70.
- Makarchenko, E. A. i Makarchenko, M. A. 2009a. Novye nakhodki (Diptera, Chironomidae, Orthoclaadiinae) na Dal'nem Vostoke i sopedel'nykh territoriyakh. VII. *Bryophaenocladus* Thienemann. (New records of chironomids (Diptera, Chironomidae, Orthoclaadiinae) in Far East and bordering territories. VII. *Bryophaenocladus* Thienemann.) - *Evraziat. ent. Zh. (= Euroas. ent. J.)* 8, *Suppl. 1*: 51-63.
- Makarchenko, E. A. i Makarchenko, M. A. 2009b. Pervye svedeniya po faune khironomid (Diptera, Chironomidae) basseina reki Langry (Severno-Zapadnyi Sakhalin). (First data on the chironomid fauna (Diptera, Chironomidae) of the Langry River Basin (North-West part of Sakhalin Island).) - *Evraziat. ent. Zh. (= Euroas. ent. J.)* 8, *Suppl. 1*: 71-77.
- Makarchenko, E. A. and Makarchenko, M. A. 2009c. *Prospilocerus amurensis* sp. n. (Diptera, Chironomidae, Orthoclaadiinae) from Amur River basin (Russian Far East). - *Evraziat. ent. Zh. (= Euroas. ent. J.)* 8: 261-263.
- Makarchenko, E. A. and Makarchenko, M. A. 2009d. New findings of chironomids (Diptera, Chironomidae) in Far East and bordering territories. IV. Subfamily Orthoclaadiinae. - *Evraziat. ent. Zh. (= Euroas. ent. J.)* 8, *Suppl. 1*: 117-124.
- Makarchenko, E. A., Makarchenko, M. A. i Yavorskaya, N. M. 2009a. Novye nakhodki khironomid (Diptera, Chironomidae) na Dal'nem Vostoke i sopedel'nykh territoriyakh. VI. *Hydrobaenus* Fries. (New records of chironomids (Diptera, Chironomidae) in Far East and bordering territories. VI. *Hydrobaenus* Fries.) - *Evraziat. ent. Zh. (= Euroas. ent. J.)* 8, *Suppl. 1*: 33-50.
- Marziali, L. 2009a. Benthic macroinvertebrates as water quality indicators in Italian lakes. - In: Hughes, S. J., Khadem, M. and Carvalho, M. A. (eds.): *Proc. 16th Int. Chironomid Symp., Bolm Mus. munic. Funchal (Hist. nat.), Suppl. 13*: 51-59.
- Masese, F. O., Muchiri, M. and Raburu, P. O. 2009a. Macroinvertebrate assemblages as biological indicators of water quality in the Moiben River, Kenya. - *Afr. J. aquat. Sci.* 34: 15-26.
- Masese, F. O., Raburu, P. O. and Muchiri, M. 2009a. A preliminary benthic macroinvertebrate index of biotic integrity (B-IBI) for monitoring the Moiben River, Lake Victoria Basin, Kenya. - *Afr. J. aquat. Sci.* 34: 1-14.
- Massaferro, J., Donato, M. and Brooks, S. J. 2009a. New placement for *Spaniotoma* (*Eukiefferiella*) *claviculata* Edwards, 1931 (Diptera: Chironomidae). - *Zootaxa* 2125: 67-68.
- Massaferro, J. I., Moreno, P. I., Denton, G. H., Vandergoes, M. and Dieffenbacher-Krall, A. 2009a. Chironomid and pollen evidence for climate fluctuations during the Last Glacial Termination in NW Patagonia. - *Quatern. Sci. Rev.* 28: 517-525.
- McGee, B. L., Pinkney, A. E., Velinsky, D. J., Ashley, J. T. F., Fisher, D. J., Ferrington, L. C. and Norberg-King, T. J. 2009a. Using the Sediment Quality Triad to characterize baseline conditions in the Anacostia River, Washington, DC, USA. - *Envir. Monit. Assess.* 156: 51-67.
- McGoff, E. and Irvine, K. 2009a. A test of the association between Lake Habitat Quality Assessment and macroinvertebrate community structure. - *Aquat. Conserv. mar. Freshwat. Ecosyst.* 19: 520-533.
- McLachlan, A. J. and Ladle, R. J. 2009a. The evolutionary ecology of detritus feeding in the larvae of freshwater Diptera. - *Biol. Rev.* 84: 133-141.
- Meltser, N., Kashi, Y. and Broza, M. 2009a. Does polarized light guide chironomids to navigate toward water surfaces? - In: Hughes, S. J., Khadem, M. and Carvalho, M. A. (eds.): *Proc. 16th Int. Chironomid Symp., Bolm Mus. munic. Funchal (Hist. nat.), Suppl. 13*: 141-149.
- Mendes, H. F. and Andersen, T. 2009a. Neotropical *Nilothauma* Kieffer, 1921, with the description of thirteen new species (Diptera: Chironomidae). - *Zootaxa* 2063: 1-45.
- Michailova, P., Ilkova, J., Hankeln, T., Schmidt, E. R., Selvaggi, A., Zampicinini, G. and Sella, G. 2009a. Somatic breakpoints, distribution of repetitive DNA and non-LTR retrotransposon insertion sites in the chromosomes of *Chironomus piger* Strenzke (Diptera, Chironomidae). - *Genetica* 135: 137-148.
- Michailova, P., Ilkova, J., Kerr, R. and White, K. 2009a. The effect of trace metals at acid and neutral conditions on the structure and

- function of polytene chromosomes in *Chironomus* species. - In: Hughes, S. J., Khadem, M. and Carvalho, M. A. (eds.): *Proc. 16th Int. Chironomid Symp., Bolm Mus. munic. Funchal (Hist. nat.), Suppl. 13*: 61-68.
- Michailova, P., Ilkova, J., Kerr, R. and White, K. 2009b. Chromosome variability in *Chironomus acidophilus* Keyl, 1960 from the Afon Goch, UK - a river subject to long-term trace metal pollution. - *Aquat. Insects* 31: 213-225.
- Michailova, P., Krastanov, B., Hankeln, T., Schmidt, E. and Kraemer, C. 2009a. *In situ* localization of the evolutionary conserved Cpy/Cty gene in the subfamily Chironominae (Chironomidae, Diptera): establishment of chromosomal homologies. - *J. zool. Syst. evol. Res.* 47: 298-301.
- Michailova, P., Szarek-Gwiazda, E. and Kownacki, A. 2009a. Effect of contaminants on the genome of some species of genus *Chironomus* (Chironomidae, Diptera) live in sediments of Dunajec River and Czorsztyn Reservoir. - *Wat. Air Soil Pollut.* 202: 245-258.
- Michelutti, N., Keatley, B. E., Brimble, S., Blais, J. M., Liu, H., Douglas, M. S. V., Mallory, M. L., Macdonald, R. W. and Smol, J. P. 2009a. Seabird-driven shifts in Arctic pond ecosystems. - *Proc. R. Soc. B* 276: 591-596.
- Mihaich, E. M., Friederich, U., Caspers, N., Hall, A. T., Klecka, G. M., Dimond, S. S., Staples, C. A., Ortego, L. S. and Hentges, S. G. 2009a. Acute and chronic toxicity testing of bisphenol A with aquatic invertebrates and plants. - *Ecotoxic. envir. Saf.* 72: 1392-1399.
- Millet, L., Arnaud, F., Heiri, O., Magny, M., Verneaux, V. and Desmet, M. 2009a. Late-Holocene summer temperature reconstruction from chironomid assemblages of Lake Anterne, northern French Alps. - *Holocene* 19: 317-328.
- Miserendino, M. L. 2009a. Effects of flow regulation, basin characteristics and land-use on macroinvertebrate communities in a large arid Patagonian river. - *Biodivers. Conserv.* 18: 1921-1943.
- Miyasaka, H. and Genkai-Kato, M. 2009a. Shift between carnivory and omnivory in stream stonefly predators. - *Ecol. Res.* 24: 11-19.
- Moiseenko, T. I., Sharov, A. N., Vandish, O. I., Kudryavtseva, L. P., Gashkina, N. A. and Rose, C. 2009a. Long-term modification of Arctic lake ecosystems: Reference condition, degradation under toxic impacts and recovery (case study Imandra Lakes, Russia). - *Limnologica* 39: 1-13.
- Moller Pillot, H. K. M. 2009a. *Chironomidae larvae of the Netherlands and adjacent lowlands – Biology and ecology of the Chironomini*. - KNNV Publ., Zeist. 270 pp.
- Móra, A. 2009a. Faunistic particularities of the chironomid assemblages (Diptera: Chironomidae) of the Upper-Tisza, NE Hungary. - In: Hughes, S. J., Khadem, M. and Carvalho, M. A. (eds.): *Proc. 16th Int. Chironomid Symp., Bolm Mus. munic. Funchal (Hist. nat.), Suppl. 13*: 5-12.
- Muscatello, J. R. and Liber, K. 2009a. Accumulation and chronic toxicity of uranium over different life stages of the aquatic invertebrate *Chironomus tentans*. - *Archs envir. Contam. Toxic.* 57: 531-539.
- Nalepa, T. F., Pothoven, S. A. and Fanslow, D. L. 2009a. Recent changes in benthic macroinvertebrate populations in Lake Huron and impact on the diet of lake whitefish (*coregonus* [sic!] *clupeaformis*). - *Aquat. Ecosyst. Hlth Mgmt* 12: 2-10.
- Nardi, J. B., Miller, L. A., Bee, C. M., Lee Jr., R. E. and Denlinger, D. L. 2009a. The larval alimentary canal of the Antarctic insect, *Belgica antarctica*. - *Arthropod Struct. Dev.* 38: 377-389.
- Neumann, D. and Woermann, D. 2009a. Physical conditions for trapping air by a microtrichia-covered insect cuticle during temporary submersion. - *Naturwissenschaften* 96: 933-941.
- Niitsuma, H. and Kato, H. 2009a. A synonymic note on *Rheopelopia joganflava* (Diptera, Chironomidae). - *Jap. J. syst. Ent.* 15: 255-257.
- Niitsuma, H. and Watson Jr, C. N. 2009a. *Bilyjomyia*, a new genus of the tribe Macropelopiini from the Holarctic (Diptera: Chironomidae). - *Zootaxa* 2166: 57-68.
- Nin, C. S., Ruppenthal, E. L. e Rodrigues, G. G. 2009a. Produção de folhico e fauna associada de macroinvertebrados aquáticos em curso d'água de cabeceira em Floresta Ombrófila do Estado do Rio Grande do Sul, Brasil. - *Acta Scient. biol. Sci.* 31: 263-271.
- Nogaro, G., Mermillod-Blondin, F., Valett, M. H., François-Carcaillet, F., Gaudet, J.-P., Lafont, M. and Gibert, J. 2009a. Ecosystem

- engineering at the sediment-water interface: bioturbation and consumer-substrate interaction. - *Oecologia* 161: 125-138.
- Nowak, C., Vogt, C., Pfenninger, M., Schwenk, K., Oehlmann, J., Streit, B. and Oetken, M. 2009a. Rapid genetic erosion in pollutant-exposed experimental chironomid populations. - *Envir. Pollut.* 157: 881-886.
- Oberholster, P. J., Botha, A.-M. and Ashton, P. J. 2009a. The influence of a toxic cyanobacterial bloom and water hydrology on algal populations and macroinvertebrate abundance in the upper littoral zone of Lake Krugersdrift, South Africa. - *Ecotoxicology* 18: 34-46.
- Obolewski, K. 2009a. Wykorzystanie makrozoobentosu do oceny stanu ekologicznego estuariowego jeziora Jamno. (Using macrozoobenthos to assess the ecological condition of the estuary Lake Jamno.) - *Ochrona środowiska* 31: 17-24.
- O'Brien, T. P., Roseman, E. F., Kiley, C. S. and Schaeffer, J. S. 2009a. Fall diet and bathymetric distribution of deepwater sculpin (*Myoxocephalus thompsonii*) in Lake Huron. - *J. Great Lakes Res.* 35: 464-472.
- Olsson, K. and Nyström, P. 2009a. Non-interactive effects of habitat complexity and adult crayfish on survival and growth of juvenile crayfish (*Pacifastacus leniusculus*). - *Freshwat. Biol.* 54: 35-46.
- Orendt, C., Wolfram, G. and Schmitt-Jansen, M. 2009a. The chironomid communities (pupal exuviae) of groyne fields in a large lowland river in Central Europe (Elbe, Germany) and their potential use for ecotoxicological field studies. - In: Hughes, S. J., Khadem, M. and Carvalho, M. A. (eds.): *Proc. 16th Int. Chironomid Symp., Bolm Mus. munic. Funchal (Hist. nat.)*, Suppl. 13: 69-75.
- Padial, A. A., Thomaz, S. M. and Agostinho, A. A. 2009a. Effects of structural heterogeneity provided by the floating macrophyte *Eichhornia azurea* on the predation efficiency and habitat use of the small Neotropical fish *Moenkhausia sanctaefilomenae*. - *Hydrobiologia* 624: 161-170.
- Park, H.-H. and Lee, J.-H. 2009a. Impact of pesticide treatment on an arthropod community in the Korean rice ecosystem. - *J. Ecol. Field Biol.* 32: 19-25.
- Park, K. and Kwak, I.-S. 2009a. Alcohol dehydrogenase gene expression in *Chironomus riparius* exposed to di(2-ethylhexyl) phthalate. - *Comp. Biochem. Physiol. Pt C* 150: 361-367.
- Park, K., Bang, H. W., Park, J. and Kwak, I.-S. 2009a. Ecotoxicological multilevel-evaluation of the effects of fenbendazole exposure to *Chironomus riparius* larvae. - *Chemosphere* 77: 359-367.
- Park, S.-Y. and Choi, J. 2009a. Genotoxic effects of nonylphenol and bisphenol A exposure in aquatic biomonitoring species: freshwater crustacean, *Daphnia magna*, and aquatic midge, *Chironomus riparius*. - *Bull. envir. Contam. Toxic.* 83: 463-468.
- Pedersen, M. L. and Friberg, N. 2009a. Influence of disturbance on habitats and biological communities in lowland streams. - *Fundam. appl. Limnol.* 174: 27-41.
- Pepinelli, M., Siqueira, T., Salles, F. F. and Shimbori, E. M. 2009a. First report of Simuliidae and Chironomidae (Diptera) living on nymphs of *Lachlania* Hagen (Ephemeroptera: Oligoneuriidae) in South America. - *Biota neotrop.* 91, bn00509012009: 245-248.
- Pestana, J. L. T., Alexander, A. C., Culp, J. M., Baird, D. J., Cessna, A. J. and Soares, A. M. V. M. 2009a. Structural and functional responses of benthic invertebrates to imidacloprid in outdoor stream mesocosms. - *Envir. Pollut.* 157: 2328-2334.
- Pestana, J. L. T., Loureiro, S., Baird, D. J. and Soares, A. M. V. M. 2009a. Fear and loathing in the benthos: Responses of aquatic insect larvae to the pesticide imidacloprid in the presence of chemical signals of predation risk. - *Aquat. Toxic.* 93: 138-149.
- Pinho, L. C. de, Mendes, H. F. and Andersen, T. 2009a. A review of *Skutzia* Reiss, 1985, with the description of three new species (Diptera: Chironomidae: Chironominae). - *J. N. Am. benthol. Soc.* 28: 196-206.
- Plóciennik, M. 2009a. Three species of Chironomidae (Diptera) new for the Polish fauna. - *Lauterbornia* 67: 107-111.
- Poi de Neiff, A., Galassi, M. E. and Franceschini, M. C. 2009a. Invertebrate assemblages associated with leaf litter in three floodplain wetlands of the Paraná River. - *Wetlands* 29: 896-906.
- Polukonova, N. V., Djomin, A. G., Muge, N. S. i Shaikovich, E. V. 2009a. Sravnenie *Chironomus usenicus* i *Ch. curabilis* s vidami gruppy *plumosus* (Diptera) po genu

- mitokhondrial'noi DNK *COI* i risunku diskov politennykh khromosom. (Comparison of *Chironomus usenicus* and *Chironomus curabilis* with species of the group *plumosus* (Diptera) inferred from the mitochondrial DNA gene *COI* and by the polytene chromosome banding pattern.) - *Genetika* 45: 1029-1035. [also published as: *Russ. J. Genet.* 45: 899-905.]
- Porinchi, D. F., MacDonald, G. M. and Rolland, N. 2009a. A 2000 year midge-based paleotemperature reconstruction from the Canadian Arctic archipelago. - *J. Paleolimnol.* 41: 177-188.
- Porinchi, D., Rolland, N. and Moser, K. 2009a. Development of a chironomid-based air temperature inference model for the central Canadian Arctic. - *J. Paleolimnol.* 41: 349-368.
- Porst, G. and Irvine, K. 2009a. Distinctiveness of macroinvertebrate communities in turloughs (temporary ponds) and their response to environmental variables. - *Aquat. Conserv. mar. Freshwat. Ecosyst.* 19: 456-465.
- Pothoven, S. A., Vanderploeg, H. A., Ludsin, S. A., Höök, T. O. and Brandt, S. B. 2009a. Feeding ecology of emerald shiners and rainbow smelt in central Lake Erie. - *J. Gt Lakes Res.* 35: 190-198.
- Power, M., Lowe, R., Furey, P., Welter, J., Limm, M., Finlay, J., Bode, C., Chang, S., Goodrich, M. and Sculley, J. 2009a. Algal mats and insect emergence in rivers under Mediterranean climates: towards photogrammetric surveillance. - *Freshwat. Biol.* 54: 2101-2115.
- Pramual, P., Gomontean, B., Buasay, V., Srikhamwiang, N., Suebkar, P., Niamlek, C., Donsinphoem, Y. and Chalath-Chieo, K. 2009a. Population cytogenetics of *Chironomus circumdatus* Kieffer, 1921 (Diptera, Chironomidae) from Thailand. - *Genetica* 135: 51-57.
- Proviz, V. I. 2009a. Kariotip i khromosmnyi polimorfizm *Chironomus solitus* Linevitsh et Erbaeva, 1971 (Diptera, Chironomidae) iz Irkutskogo vodokhranilishcha. (The karyotype and chromosomal polymorphism of *Chironomus solitus* Linevitsh et Erbaeva, 1971 (Diptera, Chironomidae) from the Irkutsk Reservoir.) - *Evraziat. ent. Zh. (= Euroas. ent. J.)* 8, Suppl. 1: 15-18.
- Proviz, V. I., Bel'kov, V. M., Proviz, L. I. i Aleksandrov, V. N. 2009a. Kariotip i morfologiya *Chironomus nigrifrons* Linevitsh et Erbaeva, 1971 (Diptera, Chironomidae) iz Irkutskogo vodokhranilishcha. (The karyotype and morphology of *Chironomus nigrifrons* Linevitsh et Erbaeva, 1971 (Diptera, Chironomidae) from the Irkutsk Reservoir. - *Evraziat. ent. Zh. (= Euroas. ent. J.)* 8, Suppl. 1: 6-14.
- Prus, P. 2009a. The dependencies between the abundance of *Chironomus* f. 1. *plumosus* and bream (*Abramis brama*) net catches in a lowland reservoir (Zegrzyński Reservoir, central Poland). - *Oceanol. hydrobiol. Stud.* 38: 15-30.
- Puntí, T., Rieradevall, M. and Prat, N. 2009a. Environmental factors, spatial variation, and specific requirements of Chironomidae in Mediterranean reference streams. - *J. N. Am. benthol. Soc.* 28: 247-265.
- Qi, X., Shi, S.-D. and Wang, X.-H. 2009a. A review of *Paratendipes* Kieffer from China (Diptera: Chironomidae). - *Aquat. Insects* 31: 63-70.
- Ramseyer, U. and Marchese, M. 2009a. Leaf litter of *Erythrina crista-galli* L. (ceibo): trophic and substratum resources for benthic invertebrates in a secondary channel of the Middle Parana River. - *Limnetica* 28: 1-10.
- Raposeiro, P. M., Hughes, S. J. and Costa, A. C. 2009a. Chironomidae (Diptera: Insecta) in Oceanic islands: New records for the Azores and biogeographic notes. - *Annl. Limnol.* 45: 59-67.
- Rasic, G., Maxwell, S. A. and Keyghobadi, N. 2009a. Characterization of microsatellite loci for the pitcher plant midge, *Metriocnemus knabi* Coq. (Diptera: Chironomidae). - *Molec. Ecol. Resourc.* 9: 1388-1391.
- Rawani, A., Haldar, K. M., Ghosh, A. and Chandra, G. 2009a. Larvicidal activities of three plants against filarial vector *Culex quinquefasciatus* Say (Diptera: Culicidae). - *Parasit. Res.* 105: 1411-1417.
- Rehn, A. C. 2009a. Benthic macroinvertebrates as indicators of biological condition below hydropower dams on west slope Sierra Nevada streams, California, USA. - *River Res. Applics* 25: 208-228.
- Reinemann, S. A., Porinchi, D. F., Bloom, A. M., Mark, B. G. and Box, J. E. 2009a. A multi-proxy paleolimnological reconstruction of Holocene climate conditions in the Great

- Basin, United States. - *Quatern. Res.* 72: 347-358.
- Riley, W. D., Pawson, M. G., Quayle, V. and Ives, M. J. 2009a. The effects of stream canopy management on macroinvertebrate communities and juvenile salmonid production in a chalk stream. - *Fish. Mgmt Ecol.* 16: 100-111.
- Rodrigues, G. G., Langton, P. H. and Scharf, B. W. 2009a. The pupal exuviae of *Chironomus crassimanus* Strenzke (Diptera: Chironomidae), an acid-resistant species from Germany. - *Zootaxa* 2026: 47-52.
- Rolland, N., Larocque, I., Francus, P., Pienitz, R. and Laperrière, L. 2009a. Evidence for a warmer period during the 12th and 13th centuries AD from chironomid assemblages in Southampton Island, Nunavut, Canada. - *Quatern. Res.* 72: 27-37.
- Rolland, N., Porinchu, D. F. and Larocque, I. 2009a. The use of high-resolution gridded climate data in the development of chironomid-based inference models from remote areas. - *J. Paleolimnol.* 41: 343-348.
- Rufer, M. M. and Ferrington Jr., L. C. 2009a. Sampling frequency required for chironomid community resolution in urban lakes with contrasting trophic states. - In: Hughes, S. J., Khadem, M. and Carvalho, M. A. (eds.): *Proc. 16th Int. Chironomid Symp., Bolm Mus. munic. Funchal (Hist. nat.), Suppl. 13*: 77-84.
- Rundio, D. E. 2009a. Community-habitat relationships in coastal streams in Big Sur, California, USA: travertine influences macroinvertebrate abundance and community structure. - *Hydrobiologia* 620: 91-108.
- Sæther, O. A. 2009a. *Telmatogeton murrayi* sp. n. from Iceland and *T. japonicus* Tokunaga from Madeira (Diptera: Chironomidae). - *Aquat. Insects* 31: 31-44.
- Sæther, O. A. 2009b. *Cryptochironomus* Kieffer from Lake Winnipeg, Canada, with a review of Nearctic species (Diptera: Chironomidae). - *Zootaxa* 2208: 1-24.
- Sánchez-Argüello, P., Fernández, C. and Tarazona, J. V. 2009a. Assessing the effects of fluoxetine on *Physa acuta* (Gastropoda, Pulmonata) and *Chironomus riparius* (Insecta, Diptera) using a two-species water-sediment test. - *Sci. tot. Envir.* 407: 1937-1946.
- Sandin, L. 2009a. The effects of catchment land-use, near-stream vegetation, and river hydromorphology on benthic macroinvertebrate communities in a south-Swedish catchment. - *Fundam. appl. Limnol.* 174: 75-87.
- Schabetsberger, R., Luger, M. S., Drozdowski, G. and Jagsch, A. 2009a. Only the small survive: monitoring long-term changes in the zooplankton community of an Alpine lake after fish introduction. - *Biol. Invasions* 11: 1335-1345.
- Schlief, J. and Mutz, M. 2009a. Effect of sudden flow reduction on the decomposition of alder leaves (*Alnus glutinosa* [L.] Gaertn.) in a temperate lowland stream: a mesocosm study. - *Hydrobiologia* 624: 205-217.
- Schmidl, J., Sulzer, P. and Kitching, R. L. 2009a. The insect assemblage in water filled tree-holes in a European temperate deciduous forest: community composition reflects structural, trophic and physicochemical factors. - *Hydrobiologia* 598: 285-303.
- Schuler, L. J., Landrum, P. F., Harwood, A. D., Tripp, E. M. and Lydy, M. J. 2009a. Joint toxicity of fluoranthene and pentachlorobenzene to *Hyalella azteca* and *Chironomus dilutus*. - *Chemosphere* 77: 399-403.
- Scott, J. J., Renaut, R. W., Buatois, L. A. and Owen, R. B. 2009a. Biogenic structures in exhumed surfaces around saline lakes: An example from Lake Bogoria, Kenya Rift Valley. - *Palaeogeogr. Palaeoclim. Palaeoecol.* 272: 176-198.
- Sealock, A. W. and Ferrington Jr., L. C. 2009a. Sampling efficiency of Chironomidae (Diptera) across disturbance gradients. - In: Hughes, S. J., Khadem, M. and Carvalho, M. A. (eds.): *Proc. 16th Int. Chironomid Symp., Bolm Mus. munic. Funchal (Hist. nat.), Suppl. 13*: 85-92.
- Sergeeva, I. V. and Anikina, M. A. 2009a. (Midges of the subfamily Tanytopodinae (Diptera, Chironomidae) of Russia: Tribes Coelotanytopodini Kieffer, 1913 and Tanytopodini Meigen, 1803.) - *Ent. Oboz.* 88: 177-184. [also published as: *Ent. Rev.* 89: 213-217.]
- Sherk, T. and Rau, G. H. 2009a. Emergence of *Ablabesmyia monilis* (Diptera: Chironomidae) from Findley Lake during warm and cold years. - In: Hughes, S. J., Khadem, M. and Carvalho, M. A. (eds.): *Proc. 16th Int. Chironomid Symp., Bolm Mus. munic. Funchal (Hist. nat.), Suppl. 13*: 199-205.

- Silver, P., Rupprecht, S. M. and Stauffer, M. F. 2009a. Temperature-dependent effects of road deicing salt on chironomid larvae. - *Wetlands* 29: 942-951.
- Skoulikidis, N. T., Karaouzas, I. and Gritzalis, K. C. 2009a. Identifying key environmental variables structuring benthic fauna for establishing a biotic typology for Greek running waters. - *Limnologica* 39: 565-66.
- Smith, B. J., Davies, P. E. and Munks, S. A. 2009a. Changes in benthic macroinvertebrate communities in upper catchment streams across a gradient of catchment forest operation history. - *For. Ecol. Mgmt* 257: 2166-2174.
- Sommer, B. and Horwitz, P. 2009a. Macroinvertebrate cycles of decline and recovery in Swan Coastal Plain (Western Australia) wetlands affected by drought-induced acidification. - *Hydrobiologia* 624: 191-203.
- Sonoda, K. C., Matthaei, C. D. and Trivinho-Strixino, S. 2009a. Contrasting land uses affect Chironomidae communities in two Brazilian rivers. - *Fundam. appl. Limnol.* 174: 173-184.
- Spies, M., Andersen, T., Epler, J. H. and Watson Jr, C. N. 2009a. Chironomidae (non-biting midges). - In: Brown, B. V., Borkent, A., Cumming, J. M., Wood, D. M., Woodley, N. E. and Zumbado, M. (eds.): *Manual of Central American Diptera, vol. 1*: 437-480. NRC Pr., Ottawa.
- Specziár, A. and Rezsú, E. T. 2009a. Feeding guilds and food resource partitioning in a lake fish assemblage: an ontogenetic approach. - *J. Fish Biol.* 75: 247-267.
- Stief, P., Poulsen, M., Nielsen, L. P., Brix, H. and Schramm, A. 2009a. Nitrous oxide emission by aquatic macrofauna. - *Proc. natn. Acad. Sci. U. S. A.* 106: 4296-4300.
- Stubbington, R., Greenwood, A. M., Wood, P. J., Armitage, P. D., Gunn, J. and Robertson, A. L. 2009a. The response of perennial and temporary headwater stream invertebrate communities to hydrological extremes. - *Hydrobiologia* 630: 299-312.
- Sun, X.-B., Wang, S.-T. and Cui, F.-Y. 2009a. (Effects of chlorine dioxide on the inactivation of the chironomid larvae.) - *Zhongguo Huanjing Kexue (China envir. Sci.)* 29: 617-621.
- Syrovátka, V. and Brabec, K. 2009a. The response of chironomid larvae to hydraulic conditions: *Synorthocladius semivirens* (DIPTERA: CHIRONOMIDAE) in two different rivers. - In: Hughes, S. J., Khadem, M. and Carvalho, M. A. (eds.): *Proc. 16th Int. Chironomid Symp., Bolm Mus. munic. Funchal (Hist. nat.), Suppl. 13*: 161-167.
- Syrovátka, V., Schenková, J. and Brabec, K. 2009a. The distribution of chironomid larvae and oligochaetes within a stony-bottomed river stretch: the role of substrate and hydraulic conditions. - *Fundam. appl. Limnol.* 174: 43-62.
- Taber, S. W. 2009a. A new species of *Polypedilum* Kieffer midge (Diptera: Chironomidae). - *SWest. Ent.* 34: 245-261.
- Takamura, N., Ito, T., Ueno, R., Ohtaka, A., Wakana, I., Nakagawa, M., Ueno, Y. and Nakajima, H. 2009a. Environmental gradients determining the distribution of benthic macroinvertebrates in Lake Takkobu, Kushiro wetland, northern Japan. - *Ecol. Res.* 24: 371-381.
- Tanikawa, T., Inoue, E. and Hirabayashi, K. 2009a. (*Limnophyes natalensis* (Kieffer, 1914) (Diptera: Chironomidae) breeding in indoor sewage.) - *Pestology* 24: 15-17.
- Taşdemir, A., Ustaoglu, M. R. ve Balık, S. 2009a. Gediz Deltası (Menemen-İzmir) Chironomidae (Diptera) Faunas'na Katkılar. (Contributions to the knowledge of Chironomidae (Diptera) fauna of the Gediz Delta (Menemen-İzmir-Turkey).) - *J. Fish. Sci.com* 3: 32-39.
- Tassou, K. T. and Schulz, R. 2009a. Effects of the insect growth regulator pyriproxyfen in a two-generation test with *Chironomus riparius*. - *Ecotoxic. envir. Saf.* 72: 1058-1062.
- Tejerina, E. G. and Paggi, A. C. 2009a. A new Neotropical species of *Oliveiriella* Wiedenbrug & Fittkau (Diptera: Chironomidae) from Argentina, with description of all its life stages. - *Aquat. Insects* 31: 91-98.
- Tellioglu, A. and Kara, T. 2009a. Seasonal abundance of Chironomidae larvae (Diptera) in Büyük Stream, Elazığ, Turkey. - *J. Freshwat. Ecol.* 24: 169-171.
- Timms, B. V. 2009a. A study of the salt lakes and salt springs of Eyre Peninsula, South Australia. - *Hydrobiologia* 626: 41-51. (+ Electr. Suppl. Mat.)

- Tixier, G., Wilson, K. P. and Williams, D. D. 2009a. Exploration of the influence of global warming on the chironomid community in a manipulated shallow groundwater system. - *Hydrobiologia* 624: 13-27.
- Tóth, M., Móra, A., Kiss, B. and Dévai, Gy. 2009a. Chironomid communities in different vegetation types in a backwater Nagy-morotva of the active floodplain of River Tisza, Hungary - In: Hughes, S. J., Khadem, M. and Carvalho, M. A. (eds.): *Proc. 16th Int. Chironomid Symp., Bolm Mus. munic. Funchal (Hist. nat.), Suppl. 13*: 169-175.
- Townsend, K. R., Pettigrove, V. J., Carew, M. E. and Hoffmann, A. A. 2009a. The effects of sediment quality on benthic macroinvertebrates in the River Murray, Australia. - *Mar. Freshwat. Res.* 60: 70-82.
- Travina, T. N. i Vvedenskaya, T. L. 2009a. Rol' khironomid (Diptera, Chironomidae) v pitanii molodi lososei i drugikh vidov ryb v reke Bolshaya. (Chironomids (Diptera, Chironomidae) in the diet of juvenile salmon and other fish in the Bolshaya River (West Kamchatka).) - *Evraziat. ent. Zh. (= Euroas. ent. J.)* 8, *Suppl. 1*: 102-108.
- Trigal, C., García-Criado, F. and Fernández-Aláez, C. 2009a. Towards a multimetric index for ecological assessment of Mediterranean flatland ponds: the use of macroinvertebrates as bioindicators. - *Hydrobiologia* 618: 109-123.
- Trivinho-Strixino, S. and Strixino, G. 2009a. A new species of *Pelomus* Reiss, 1989 (DIPTERA: CHIRONOMIDAE) from Southeastern Brazil, with the description of immature stages. - In: Hughes, S. J., Khadem, M. and Carvalho, M. A. (eds.): *Proc. 16th Int. Chironomid Symp., Bolm Mus. munic. Funchal (Hist. nat.), Suppl. 13*: 217-225.
- Tullos, D. D., Penrose, D. L., Jennings, G. D. and Cope, W. G. 2009a. Analysis of functional traits in reconfigured channels: implications for the bioassessment and disturbance of river restoration. - *J. N. Am. benthol. Soc.* 28: 80-92.
- Tyagi, A., Ryme, J., Brodin, D., Östlund Farrants, A. K. and Visa, N. 2009a. SWI/SNF associates with nascent pre-mRNPs and regulates alternative pre-mRNA processing. - *PLoS Genet.* 5, 5, e1000470: 15 pp.
- Utz, R. M., Hilderbrand, R. H. and Boward, D. M. 2009a. Identifying regional differences in threshold responses of aquatic invertebrates to land cover gradients. - *Ecol. Indicators* 9: 556-567.
- Uzarski, D. G., Burton, T. M., Kolar, R. E. and Cooper, M. J. 2009a. The ecological impacts of fragmentation and vegetation removal in Lake Huron's coastal wetlands. - *Aquat. Ecosyst. Hlth Mgmt* 12: 45-62.
- Vallenduuk, H. 2009a. Larvae of the Tanyptodinae (Diptera: Chironomidae): New (neglected) characters and the making of a new key. - *Proc. Neth. ent. Soc. Meet.* 20: 55-66.
- Varadinova, E., Pehlivanov, L., Moskova, G. and Uzunov, Y. 2009a. Development and actual state of the macrozoobenthic fauna in the Srebarna Biosphere Reserve (north-east Bulgaria). - *C. r. Acad. Bulg. Sci.* 62: 243-248.
- Vedamanikam, V. J. 2009a. Formation of resistance in the *Chironomus plumosus* to four pesticides over 45 generations. - *Toxic. envir. Chem.* 91: 187-194.
- Vedamanikam, V. J. and Shazili, N. A. M. 2009a. Observations of mouthpart deformities in the *Chironomus* larvae exposed to different concentrations of nine heavy metals. - *Toxic. envir. Chem.* 91: 57-63.
- Vedamanikam, V. J. and Shazili, N. A. M. 2009b. Comparative toxicity of nine metals to two Malaysian aquatic dipterian larvae with reference to sediment composition. - *Toxic. envir. Chem.* 91: 129-136.
- Vedamanikam, V. J. and Shazili, N. A. M. 2009c. The chironomid larval tube, a mechanism to protect the organism from environmental disturbances? - *Toxic. envir. Chem.* 91: 171-176.
- Vink, J. P. M. 2009a. The origin of speciation: Trace metal kinetics over natural water/sediment interfaces and the consequences for bioaccumulation. - *Envir. Pollut.* 157: 519-527.
- Vinogradova, E. M., Riss, H. W. and Spies, M. 2009a. New species of *Tanytarsus* van der Wulp, 1874 (Diptera: Chironomidae) from Central America. - *Aquat. Insects* 31: 11-17.
- Virtanen, R., Ilmonen, J., Paasivirta, L. and Muotka, T. 2009a. Community concordance between bryophyte and insect assemblages in boreal springs: a broad-scale study in isolated habitats. - *Freshwat. Biol.* 54: 1651-1662.

- Vonesh, J. R. and Kraus, J. M. 2009a. Pesticide alters habitat selection and aquatic community composition. - *Oecologia* 160: 379-385.
- Vonesh, J. R., Kraus, J. M., Rosenberg, J. S. and Chase, J. M. 2009a. Predator effects on aquatic community assembly: disentangling the roles of habitat selection and post-colonization processes. - *Oikos* 118: 1219-1229.
- Wang, Y. V., O'Brien, D. M., Jenson, J., Francis, D. and Wooller, M. J. 2009a. The influence of diet and water on the stable oxygen and hydrogen isotope composition of Chironomidae (Diptera) with paleoecological implications. - *Oecologia* 160: 225-233.
- Weston, D. P., Lentz, R. D., Cahn, M. D., Ogle, R. S., Rothert, A. K. and Lydy, M. J. 2009a. Toxicity of anionic polyacrylamide formulations when used for erosion control in agriculture. - *J. envir. Qual.* 38: 238-247.
- Wiedenbrug, S. and Trivinho-Strixino, S. 2009a. *Ubatubaneura*, a new genus of the *Corynoneura* group (Diptera: Chironomidae: Orthoclaadiinae) from the Brazilian Atlantic Forest. - *Zootaxa* 1993: 41-52.
- Wiedenbrug, S., Mendes, H. F., Pepinelli, M. and Trivinho-Strixino, S. 2009a. Review of the genus *Onconeura* Andersen *et* Sæther (Diptera: Chironomidae), with the description of four new species from Brazil. - *Zootaxa* 2265: 1-26.
- Wijngaarden, R. P. A. van, Barber, I. and Brock, T. C. M. 2009a. Effects of pyrethroid insecticide gamma-cyhalothrin on aquatic invertebrates in laboratory and outdoor microcosm tests. - *Ecotoxicology* 18: 211-224.
- Williams, M., Cook, E., van der Kaars, S., Barrows, T., Shulmeister, J. and Kershaw, P. 2009a. Glacial and deglacial climatic patterns in Australia and surrounding regions from 35 000 to 10 000 years ago reconstructed from terrestrial and near-shore proxy data. - *Quatern. Sci. Rev.* 28: 2398-2419.
- Wissinger, S. A., Greig, H. and McIntosh, A. 2009a. Absence of species replacements between permanent and temporary lentic communities in New Zealand. - *J. N. Am. benthol. Soc.* 28: 12-23.
- Wright, I. A. and Burgin, S. 2009a. Effects of organic and heavy metal pollution on chironomids within a pristine upland catchment. - *Hydrobiologia* 635: 15-25.
- Wuelker, W., Martin, J., Kiknadze, I. I., Sublette, J. E. and Michiels, S. 2009a. *Chironomus blaylocki* sp. n. and *C. bifurcatus* sp. n., North American species near the base of the *decorus*-group (Diptera: Chironomidae). - *Zootaxa* 2023: 28-46.
- Yang, Y., Hunter, W., Tao, S. and Gan, J. 2009a. Effects of black carbon on pyrethroid availability in sediment. - *J. agric. Food Chem.* 57: 232-238.
- Yasué, M. and Dearden, P. 2009a. The importance of supratidal habitats for wintering shorebirds and the potential impacts of shrimp aquaculture. - *Envir. Mgmt* 43: 1108-1121.
- Yavorskaya, N. M. i Makarchenko, M. A. 2009a. Pervye svedeniya po faune khironomid (Diptera, Chironomidae) Bol'shekhkhtsirskogo gosudarstvennogo prirodnogo zapovednika (Khabarovskii krai). (First data on the chironomid fauna (Diptera, Chironomidae) of the Bolshekhkhtsirski nature Reserve (Khabarovsk Territory).) - *Evrasiat. ent. Zh. (= Euroas. ent. J.)* 8, Suppl. 1: 93-96.
- Yoshimi, T., Odagiri, K., Hiroshige, Y., Yokobori, S.-i., Takahashi, Y., Sugaya, Y. and Miura, T. 2009a. Induction profile of HSP70-cognate genes by environmental pollutants in Chironomidae. - *Envir. Toxic. Pharmac.* 28: 294-301.
- Zelentsov, N. I. 2009a. Fauna khironomid (Diptera, Chironomidae) Khibinskikh ozer Kol'skogo poluostrova. (The chironomid fauna (Diptera, Chironomidae) of the Khibiny's lakes of Kola Pensinsula.) - *Evrasiat. ent. Zh. (= Euroas. ent. J.)* 8, Suppl. 1: 89-92.
- Zinchenko, T. D., Makarchenko, M. A. i Makarchenko, E. A. 2009a. Novyi vid roda *Cricotopus* van der Wulp (Diptera, Chironomidae) iz solenoi reki basseina ozera El'ton (Volgogradskaya oblast', Rossiya). (A new species of *Cricotopus* van der Wulp (Diptera, Chironomidae) from a saline river of the El'ton Lake basin (Volgograd Region, Russia).) - *Evrasiat. ent. Zh. (= Euroas. ent. J.)* 8, Suppl. 1: 83-88.
- Zorina, O. V. 2009a. *Synendotendipes shanduyensis* sp. n. (Diptera, Chironomidae, Chironominae) from elevated lakes of Sikhote-Alin' Biosphere Nature Reserve. - *Evrasiat. ent. Zh. (= Euroas. ent. J.)* 8, Suppl. 1: 78-82.

- Zorina, O. V. i Zinchenko, T. D. 2009a. Novyi vid roda *Tanytarsus* van der Wulp (Diptera, Chironomidae) iz solenoi reki basseina reki El'ton (Volgogradskaya oblast', Rossiya). (A new species of *Tanytarsus* van der Wulp (Diptera, Chironomidae) from a saline river of the El'ton Lake basin (Volgograd Region, Russia). - *Evrasiat. ent. Zh. (= Euroas. ent. J.)* 8, Suppl. 1: 105-110.
- Abstracts from the XVIIth International Symposium on Chironomidae in Nankai University, Tianjin, China, 2009** (sequence in abstracts book, Wang, X.-H., ed.)
- A. Oral presentations:
- Martin, J. 2009a. From bands to base pairs: Problems in the identification of *Chironomus* species, using the example of *Chironomus oppositus* Walker. - 11.
- Al-Shami, S. A., Che Salmahi, M. R., Siti Azizah, M. N. and Abu Hassan, A. 2009b. Distribution of Chironomidae (Diptera) under the influence of several environmental parameters in polluted rivers of Juru River Basin, Penang, Malaysia. - 12.
- Anderson, A. M., Bouchard, R. W. and Ferrington, L. C. 2009a. Hibernial emergence of Chironomidae in relation to stream size in Kansas. - 13.
- Ashe, P. and O'Connor, J. P. 2009a. Part 1 of a world catalogue of Chironomidae (Diptera). - 14.
- Brabec, K., Jarkovsky, J. and Kubosova, K. 2009a. Occurrence and co-occurrence of stream chironomids: associations with environmental gradients and community characteristics. - 15.
- Chon, T., Tang, H., Qu, X., Bae, M., Song, M. and Park, Y. 2009a. Community structure and species abundance distribution in chironomids collected in streams in different pollution levels. - 16.
- Ebrahimnezhad, M. and Allahbakhshi, E. 2009a. Chironomidae (Diptera) genera from Golpayegan River, Iran. - 17.
- Ekrem, T., Willassen, E. and Stur, E. 2009a. Molecular phylogeny of *Micropsectra* and allies: indications of new generic synonyms (Diptera: Chironomidae): - 18-19.
- Ermolaeva, O. 2009b. Comparative karyological analysis of the subfamily Diamesinae (Diptera, Chironomidae). - 20.
- Ferrington Jr., L. C., Westrick, M. and Karns, B. 2009a. Alpha, beta and gamma diversities of Chironomidae emergence from 20 spring-runs in the Saint Croix River basin, Minnesota. - 21-22.
- Golygina, V. V., Istomina, A. G. and Kiknadze, I. I. 2009a. Interspecific hybridization between sibling species of *Chironomus plumosus* group (Diptera: Chironomidae). - 23-24.
- Gresens, S. E. 2009a. Response of Chironomidae to multiple gradients of urban impact. - 25.
- Gunderina, L. I. and Katokhin, A. V. 2009a. Variation and divergence of rDNA ITS-1 region in species of genus *Chironomus* (Diptera: Chironomidae). - 26-27.
- Inoue, E., Tsukada, M. and Hirabayashi, K. 2009a. Structural analyses of silk and nesting tubes of chironomid larvae (Diptera). - 28.
- Kiknadze, I. and Istomina, A. 2009a. Karyological study of *Chironomus* species from the Netherlands. - 29-30.
- Kaga, K., Kasuya, S., Kobayashi, T. and Ohtaka, A. 2009a. Identification of chironomid species by DNA sequence - especially genus *Hydrobaenus* including *H. sp.* Tsugaru. - 31.
- Kobayashi, T., Ohtaka, A., Kasuya, S. and Kaga, K. 2009a. *Hydrobaenus* sp. Tsugaru with extremely short male antennae from Tsugaru Peninsula, northern Japan (Diptera: Chironomidae). - 32-33.
- Langton, P. H. 2009a. Pupal exuviae structure further elaborated. - 34.
- Lencioni, V., Marziali, L. and Rossaro, B. 2009b. Diversity and distribution of chironomids in natural and impacted springs in the Italian Prealps and Alps. - 35.
- Liu, Y., Ji, C. and Chon, T. 2009a. Automatic detection of response behaviors of *Chironomus riparius* after exposure to toxic chemicals by using computational methods. - 36.
- Makarchenko, E. A. and Makarchenko, M. A. 2009e. Fauna and taxonomy of the Orthocladiinae (Diptera: Chironomidae) of the Russian Far East. - 37.
- Marziali, L., Rossaro, B. and Buffagni, A. 2009a. Temporal and spatial variability of chironomid assemblages in a large lowland river (Po River, Northern Italy). - 38-39.
- Rossaro, B., Marziali, L., Boggero, A. and Lencioni, V. 2009a. Benthic

- macroinvertebrates (Chironomidae, Oligochaeta) in Italian lakes. - 40-41.
- McBean, J. E. 2009a. *Microtendipes* Kieffer prohumeral reversed setae – gratis mutation or functional asset? - 42.
- McBean, S. F. 2009a. Questioning feeding related factors in niche separation in chironomid larvae, in particular mentum profile of Orthoclaadiinae and Chironominae. - 43.
- Nath, B. B. and Babrekar, A. A. 2009a. Implications of hypoxic stress in *Chironomus ramosus* larvae with reference to iron. - 44.
- Przhiboro, A. and Sæther, O. A. 2009a. Littoral chironomid communities of two small lakes in Northern Karelia (Russia) studied by emergence traps: species composition and changes over a 30-years period. - 45-46.
- Stur, E. and Ekrem, T. 2009a. Exploring unknown life stages of Arctic Chironomidae (Diptera) with DNA barcoding. - 47.
- Yamamoto, M. and Yamamoto, N. 2009a. A review of *Yaeprius isigaabeus* Sasa & Susuki, 2000 (Diptera: Chironomidae). - 48.
- Yamamoto, N., Hirowatari, T. and Yamamoto, M. 2009a. The subgenus *Pentapedilum* of the genus *Polypedilum* (Diptera: Chironomidae) in Iriomote Island, the Ryukyus, Japan. - 49.
- B. Poster presentations:**
- Ali, A., Leckel Jr., R. J. and Duxbury, C. V. 2009a. Chironomidae investigations in selected Florida wetlands. - 51-52.
- Andersen, T., Mendes, H. F., Pinho, L. C., Roque, F. O. and Spies, M. 2009a. An updated checklist of Brazilian Chironomidae (Diptera). - 53-54.
- Ashe, P. 2009a. A bibliography, with dates of publication, of J. J. Kieffer's publications on Chironomidae (Diptera). - 55.
- Caldwell, B. A. 2009b. A novel pupa of *Pseudorthocladus* Goetghebuer, new Nearctic record for *Doithrix villosa* Sæther & Sublette, and variation in *Doithrix dillonae* Cranston & Oliver (Diptera: Chironomidae). - 56.
- Cheng, M. and Wang, X. 2009b. *Ablabesmyia* Johannsen (Diptera: Chironomidae: Tanytopodinae) from China. - 57.
- Przhiboro, A., Ekrem, T. and Stur, E. 2009a. Bionomics and taxonomy of *Tanytarsus recurvatus* Brundin, 1947. - 58.
- Ermolaeva, O. 2009c. Comparative karyological analysis of the subfamily Prodiamesinae (Diptera, Chironomidae). - 59.
- Fu, Y., Andersen, T. and Mendes, H. F. 2009a. *Paratrachocladus* Santos Abreu from Patagonia (Diptera: Chironomidae: Orthoclaadiinae). - 60.
- Fu, Z., Inoue, E., K. Yoshizawa, K., Yoshida, N., Kazama, F. and K. Hirabayashi, K. 2009a. Horizontal distribution of chironomid larvae (Diptera) in Lake Saiko, Japan. - 61.
- Grzybkowska, M. and Głowacki, Ł. 2009a. Chironomidae (Diptera) diversity in lowland rivers of different orders in central Poland. - 62.
- Hirabayashi, K., Yoshizawa, K., Oga, K., Yoshida, N., Ariizumi, K. and Azama, F. K. 2009a. Population dynamics of benthic macroinvertebrates in a eutrophic-mesotrophic lake, Lake Kawaguchi, Japan. - 63-64.
- Hu, Y. 2009a. A case study for effects of bt rice on non-biting midges in paddy field. - 65.
- Ikutama, E., Inoue, E. and Hirabayashi, K. 2009a. Density dynamics of chironomid larvae (Diptera) in the middle reaches of the Shinano River, Japan: the response to floods. - 66.
- Kawai, K., Okamoto, H. and Imabayashi, H. 2009a. A fundamental study on range expansion of chironomids. - 67.
- Lencioni, V. and Rossaro, B. 2009a. Chironomids of springs, rivers and lakes in five provinces and in the Beijing Municipality of the People's Republic of China. - 68.
- Bernabò, P., Caputi, L., Frassanito, R., Guella, G., Jousson, O. and Lencioni, V. 2009a. Cold resistance in *Pseudodiamesa branickii* (Nowicki): physiological, molecular and biochemical aspects. - 69.
- Majumdar, U. 2009a. Indian species of *Glyptotendipes* Kieffer (Diptera: Chironomidae) with a short account of some aspects of biology. - 70.
- Makarchenko, E. A. 2009a. *Pseudodiamesa nivosa* (Goetghebuer, 1928) is not synonym of *Pseudodiamesa arctica* (Malloch, 1919) (Diptera: Chironomidae). - 71.
- Marziali, L., Rossaro, B. and Lencioni, V. 2009a. Chironomid drift behavior (Insecta, Diptera) in an arctic stream (Svalbard Islands) under natural and induced light conditions. - 72-73.

- Wiedenbrug, S., Mendes, H. F. and Trivinho-Strixino, S. 2009a. New species of the genus *Onconeura* Andersen & Sæther, 2005 (Diptera, Chironomidae) from Brazil. - 74.
- Móra, A. 2009a. The diversity of chironomid assemblages of streams in Lake Balaton's catchment area. - 75.
- Pinho, L. C. de and Froehlich, C. G. 2009a. Three new Neotropical species of *Fissimentum* Cranston *et* Nolte, 1996 (Diptera: Chironomidae: Chironominae). - 76.
- Plank, A. 2009a. Chip – chironomid identification program: build your own determination key. - 77-78.
- Raposeiro, P., Costa, A. C. and Hughes, S. J. 2009a. Preliminary results of the use of chironomid pupal exuviae technique in oceanic islands: Azores Archipelago. - 79-80.
- Sergeeva, I. V. 2009a. Fauna and ecology of the Russia Tanypodinae (Diptera, Chironomidae). - 81.
- Shimodoi, T., Kawai, K. and Imabayashi, H. 2009a. Fundamental study of fish effectiveness to control chironomid mass emergence. - 82.
- Sonoda, K. C., Ortega, E., Vettorazzi, C. A. and Trivinho-Strixino, S. 2009a. Chironomidae assemblage from four Brazilian watersheds under different anthropogenic pressure. - 83-84.
- Tóth, M., Kőműves, M. and Dévai, G. 2009a. Spatio-temporal distribution of phytal-dwelling chironomid assemblages (Diptera: Chironomidae) in submerged vegetation of two Hungarian backwaters. - 85.
- Watanabe, T., Kawai, K. and Imabayashi, H. 2009a. A fundamental study to control adult chironomid midges by using the light traps. - 86.
- Yan, C. and Wang, X. 2009a. A taxonomic discussion on a valid species of *Harnischia longispuria* Wang & Zheng, 1993 (Diptera: Chironomidae). - 87.
- Zinchenko, T. D. and Golovatyuk, L. V. 2009a. Responds of the river chironomid communities (inflows of the hypersaline lake Elton, the Low Volga, south of Russia) to high mineralization. - 88-89.
- C. Abstracts of Non-attending Colleagues
- (Note: Several authors of contributions in paragraphs A and B did not attend the meeting either)
- Chavan, R. J. 2009a. Diversity of chironomids (Diptera) in urban lakes at Aurangabad, India. - 91.
- Jacobsen, R. E. 2009b. *Tanytarsus rexilius*, new species, from oligotrophic marshes in the Florida Everglades. - 92.
- Morozova, Y. Y. 2009a. Ecologically-morphological analysis of *Cryptochironomus* Kieffer (Diptera: Chironomidae) species of the Volga River. - 93.
- Zhang, E. 2009a. Fossil midges (Diptera: Chironomidae) as paleoecological indicators in Chinese lakes. - 94.